

# Risk factors for obstetric anal sphincter injuries during vaginal delivery: can we reduce the burden?

Irene Porcari<sup>1</sup>, Simone Garzon<sup>1,\*</sup>, Sara Loreti<sup>1</sup>, Mariachiara Bosco<sup>1</sup>, Rossana Di Paola<sup>1</sup>, Paola Pomini<sup>1</sup>, Ricciarda Raffaelli<sup>1</sup>, Fulvio Leopardi<sup>2</sup>, Stefano Uccella<sup>1</sup>, Massimo Franchi<sup>1</sup>, Pier Carlo Zorzato<sup>1</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, University of Verona, 37126 Verona, Italy

<sup>2</sup>Department of Surgery, University of Verona, 37126 Verona, Italy

\*Correspondence: [simone.garzon@univr.it](mailto:simone.garzon@univr.it) (Simone Garzon)

DOI: [10.31083/j.ceog4806201](https://doi.org/10.31083/j.ceog4806201)

This is an open access article under the CC BY 4.0 license (<https://creativecommons.org/licenses/by/4.0/>).

Submitted: 13 August 2021 Revised: 7 September 2021 Accepted: 21 October 2021 Published: 15 December 2021

**Background:** Third- and fourth-degree perineal tears are associated with significant discomfort and impact on women's quality of life after labor. We reviewed the literature on risk factors for obstetric anal sphincter injuries (OASIS), focusing on modifiable risk factors for OASIS to help obstetricians prevent them. **Methods:** We searched MEDLINE, EMBASE, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews, and Scopus using key search terms. We selected clinical studies, systematic reviews, and meta-analyses in English investigating antepartum and intrapartum factors associated with OASIS. Three researchers independently selected studies and documented outcomes. **Results:** We identified forty-two relevant articles for inclusion. Among antepartum factors, primiparity, neonatal birth weight, and ethnicity were associated with a higher risk of OASIS. Instrumental delivery, midline episiotomy, duration of the second stage of labor, persistent occiput posterior position, and labor augmentation were those intrapartum factors reported associated with OASIS. **Conclusions:** Multiple ante- and intrapartum factors are associated with a higher risk of OASIS. The actual modifiable factor is episiotomy during the second stage of labor. However, literature reporting episiotomy associated with a reduction in OASIS prevalence during instrumental delivery is limited. These results may help obstetricians recognize women at higher risk of developing severe perineal tears and recommend further investigation on the role of episiotomy in an operative delivery.

## Keywords

Obstetric anal sphincter injury; OASIS; Perineal tears; Obstetric trauma; Episiotomy

## 1. Introduction

Perineal tears are caused by the straining of the pelvic floor structure during vaginal delivery. As severity concerns, they are classified into four levels which reflect the involved anatomic structures [1]. Third- and fourth-degree lacerations are known as severe obstetric anal sphincter injuries (OASIS), and they include injuries that involve the anal sphincter only (third-degree) and those extending to the anal mucosa (fourth-degree).

The incidence of OASIS represents a maternal wellbeing indicator, and it should not exceed 5% of all perineal tears in labor [2]. In the United States, the rate of third-degree lac-

eration is 3.3%, and of fourth-degree laceration is 1.1% [3]. However, almost 20% of OASIS are estimated to be missed and thus unrepaired during the postpartum period with associated long-term complications [4].

OASIS results in short-term morbidity such as pain, infection, hemorrhage [5], and long-term sequelae like dyspareunia and urinary and anal incontinence [6, 7]. Stool and flatus incontinence occurs in up to 50% of women after OASIS [8]. These morbidities profoundly impact women's psychological wellbeing and are associated with a high risk of postpartum depression in women with lower resilience [9]. Moreover, the associated surgical repair is at high risk of needlestick injury among obstetricians [10, 11]. Nevertheless, prevention strategies for severe perineal lacerations are still far from being established. Although it represents an essential topic for obstetricians, inadequate evidence is available in the literature due to the heterogeneity of studies' conclusions and the absence of randomized trials.

In this scenario, defining risk factors for perineal trauma after vaginal delivery would be helpful to identify women at risk for developing OASIS and find out possible preventive measures. Therefore, this review aims to analyze risk factors associated with OASIS and identify women at higher risk for severe perineal tears. This is of primary importance to help obstetricians understand how to reduce the incidence and impact of these events, particularly whether modifiable risk factors are identified.

## 2. Materials and methods

For this review, a comprehensive search of several databases was conducted from inception up to May 2021. The searched databases were MEDLINE, Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Daily, Ovid EMBASE, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews, and Scopus.

The research strategy included the combinations of the following medical terms: "perineal lacerations; perineal trauma; perineal tear; childbirth trauma; birth trauma; ob-

stetric trauma; obstetric anal sphincter injury; OASIS; third-degree tear; fourth-degree tear". We selected clinical studies, systematic reviews, and meta-analyses in English language investigating antepartum and intrapartum factors associated with OASIS. We selected studies and systematic reviews or meta-analyses of studies, including pregnant women who delivered vaginally and reporting data regarding the occurrence of OASIS and intra or antepartum factors associated with OASIS occurrence. No additional inclusion or exclusion criteria were used. No specific interventions were investigated.

We did not use a systematic approach in reporting results due to the heterogeneous and extensive amount of literature on this topic; therefore, we conducted a narrative review and reported the most relevant studies to provide the reader with a complete and synthetic overview of the risk factors for severe perineal tears to prevent perineal lesions during delivery. Data are presented as absolute numbers (percentage). No statistical analysis was performed.

### 3. Results

Forty-two articles have been selected for inclusion in the present review. From the included studies, we identified multiple modifiable and non-modifiable factors associated with OASIS.

#### 3.1 Antenatal factors

Pergialiotis *et al.* [12] found that primiparous women have the highest rate of severe perineal lacerations (RR 1.59, [95% confidence interval (CI), 1.45–1.75];  $p < 0.001$ ). In a cohort of 2754 women, Smith *et al.* [13] reported an overall OASIS rate of 6.6% in nulliparous women and 2.7% in multiparous ones.

Asian women were reported to have a higher rate of severe perineal tears than Caucasian women. Extensive literature observed this association; however, the etiological factors are still unclear. Differences in pelvic floor anatomy and function between racial groups have been proposed as determinants of this higher susceptibility of women belonging to specific races [14, 15].

#### 3.2 Intrapartum factors

Some authors found an increased rate of OASIS after births of macrosomic infants [16–19]. In a study of 3356 women who delivered infants with a birth weight greater than 4500 g, 4.4% of them suffered from severe perineal trauma. This rate was higher compared with women who had a non-macrosomic fetus [17]. Stotland *et al.* [18] included 146526 births and calculated the rate of fourth-degree lacerations: 1.5% when the neonatal birth weight was less than 4000 g, 2.8% when the neonatal weight was between 4000 g and 4499 g, 4.5% for neonatal birth weight below 4500 g, and 7% for infants who weighted 5000 g. Conversely, other authors did not find an association between severe perineal trauma and birth weight in macrosomic fetuses [20, 21]. A retrospective study on 34685 term singleton pregnancies, of which 2077 were characterized by a fetal birth weight greater than 4000 g, showed no difference in OASIS rate between

macrosomic and non-macrosomic fetuses [20]. In a meta-analysis of 716031 patients, neonatal birth weight was significantly higher in women who developed severe perineal lacerations than in the control group. However, the mean difference was 163.71 g. The weight difference between the two groups was so slight that clinical relevance has been questioned. Therefore, the importance of fetal birth weight on the OASIS rate is still debated [12].

Mazouni *et al.* [22] observed no correlation between the obstetric maneuvers used to solve fetal shoulder dystocia (SD) and perineal lesions. Conversely, Gauthaman *et al.* [23] reported a higher risk of OASIS when internal maneuvers were used for SD management than those women who underwent eutocic vaginal birth (OR 2.18, [95% CI, 1.17–4.0]). This is particularly true when four or more maneuvers are used. In a multivariate analysis, internal maneuvers such as Woods' and the reverse Woods' screw maneuvers were found to be significant risk factors, while delivering the posterior fetal arm at first is not a risk factor for OASIS (OR 1.74, [95% CI, 0.84–3.6];  $p = 0.135$ ) [23].

Several authors observed that the rate of third and fourth-degree perineal tears was associated with the length of the second labor stage [24, 25]. This is true for both nulliparous and multiparous women [26]. Ramm *et al.* [27], in a retrospective study, compared women whose second stage of labor lasted at least 180 minutes to those with the second stage of labor that lasted less than 60 minutes, describing a three times higher OASIS rate in the first group (adjusted OR 3.20, [95% CI, 2.62–3.89]). Consistently, labor augmentation is an independent risk factor for the development of severe lacerations (RR 1.46, [95% CI, 1.32–1.62]) [12, 28, 29].

Conversely, labor induction was not associated with severe perineal tears. The ARRIVE trial compared elective labor induction at 39 weeks in low-risk nulliparous women with expectant management. They demonstrated that induction results in a significantly lower frequency of cesarean delivery [30]. In a secondary analysis of the ARRIVE trial, the OASIS rate was similar between women with induced labor and those managed expectantly [31]. The same results were reported in a meta-analysis, showing that labor induction had no significant effect on the OASIS risk (RR 1.05, [95% CI, 0.97–1.15], 716031 women) [12].

Conflicting results have been reported regarding the association between the use of epidural analgesia and OASIS. Some studies suggested that it may protect against severe perineal tears [13]. Loewenberg-Weisband *et al.* [32], in the first analysis, found a significant association between the use of epidural analgesia and OASIS. However, after adjusting the results with the effects of parity, this association disappeared (OR 0.95, [95% CI, 0.69–1.29]). Still, other studies reported epidural analgesia as a risk factor for OASIS (RR 1.21, [95% CI 1.08–1.36], 716,031 women) [12].

Occiput posterior position at delivery is an independent risk factor for severe lacerations [33] and is associated with a significantly lower rate of spontaneous vaginal births [12,

34]. One possible explanation might be related to the fetal head's inability to flex onto the chest, resulting in an enlargement of the fetal head diameter at the pelvic outlet [35]. Moreover, occiput posterior position is associated with prolonged first and second stages of labor and increased medical interventions such as oxytocin augmentation and assisted delivery [34].

Episiotomy is a surgical procedure that consists of vaginal orifice enlargement by incising the perineum to facilitate fetal delivery. Its role as a preventive maneuver for OASIS or risk factors has become a highly debated topic in the last decade.

Episiotomy can be performed at the end of the second stage of labor when the perineum is seen as an obstacle to the completion of fetal birth [36]. Different types of incisions were described in the literature, such as the midline, mediolateral, modified-median, J-shaped, and lateral ones [37, 38]. However, the two most used incisions are the midline (median) and the mediolateral ones. The former is more frequently performed in the United States, while the latter is more used in Europe [38]. The literature agrees that midline episiotomy is associated with an increased risk of OASIS as compared with mediolateral episiotomy [12, 27], and International Guidelines, such as the American College of Obstetricians and Gynecologists (ACOG) and Royal College of Obstetricians and Gynecologists (RCOG), discourage its use and agree that mediolateral episiotomy may be preferable [39, 40].

In women undergoing a non-operative vaginal delivery, mediolateral episiotomy was assumed to be protective against severe perineal tears for a long time. Nevertheless, it is well demonstrated that routine use of episiotomy during eutocic spontaneous vaginal delivery is not justified in reducing perineal trauma [41, 42]. According to Cochrane database review, the selective use of episiotomy compared with the routine one resulted in a 30% decrease of severe perineal trauma (RR 0.70, [95% CI, 0.52–0.94]; 5375 women) [43]; this was only applicable to trials where the difference in episiotomy rates between the two groups was more than 30%. Moreover, in a meta-analysis of 2020, the authors demonstrated that mediolateral episiotomy tends to increase the risk of severe perineal lacerations. However, the overall effect does not reach statistical significance ( $p = 0.08$ ) [12].

A different situation is the operative vaginal delivery, which has been recognized as a major risk factor for OASIS, particularly when other risk factors coexist [27]. In the literature, forceps for operative vaginal delivery was associated with an increased risk of OASIS [44]. Similarly, many studies reported even vacuum extraction as a significant risk factor for perineal trauma [12, 45]. However, Ramm *et al.* [27] included a cohort of 22822 vaginal deliveries reporting a 4–6 times higher OASIS rate among patients who underwent forceps-assisted deliveries than vacuum-assisted deliveries. For this and additional evidence, the actual contribution of vacuum-assisted delivery to OASIS risk is still debated. Indeed, the actual contribution of vacuum-assisted delivery to

the occurrence of OASIS is challenging to estimate, as other determinants may confound its impact.

Levin *et al.* [46] analyzed the distribution of tear types in spontaneous vaginal births compared to vacuum-assisted deliveries. They evaluated 23272 vaginal deliveries of primiparous women. The univariate analysis reported a higher OASIS rate in the vacuum delivery group (2.3%) than the spontaneous vaginal births group (1.7%). However, when multivariate regression analysis was performed, vacuum-assisted delivery was not independently associated with an increased risk of OASIS. In addition, in the vacuum-assisted delivery group, aspects like maternal age >35 years old, weight gain during gestation, oxytocin administration, epidural anesthesia, occiput posterior fetal position, and longer second-stage duration were found more frequently.

Similarly, although many authors suggested that mediolateral episiotomy prevented OASIS during operative vaginal delivery, the role of mediolateral episiotomy during operative vaginal delivery to prevent OASIS has not been confirmed by large randomized controlled trials (RCT) or meta-analysis [47]. Historically, international guidelines did not support routine mediolateral episiotomy in operative vaginal deliveries to reduce OASIS incidence. However, recently, RCOG has stated that mediolateral episiotomy prevents OASIS in primiparous women [40]. Van Bavel *et al.* [48] evaluated the effect of mediolateral episiotomy on OASIS rate during operative vaginal delivery in a retrospective study in 2018. The incidence of OASIS in primiparous women who underwent mediolateral episiotomy was 2.5%, and in those who did not was 14%, with an episiotomy rate of 84.5%. They reported a protective effect of mediolateral episiotomy during instrumental delivery in both primiparous (adjusted OR 0.14, [95% CI, 0.13–0.15]) and multiparous (adjusted OR 0.23, [95% CI, 0.21–0.27]) women [48]. De Vogel *et al.* [49] corroborated the protective effect of episiotomy on perineal tears during operative vaginal delivery (OR 0.17, [95% CI, 0.12–0.24]). Data from 2861 operative vaginal delivery showed an OASIS rate of 3.3% in women managed with episiotomy and 15.6% in those managed without it. The global episiotomy rate was 81%.

Conversely, other authors suggested that the risk of OASIS was not affected by the use of episiotomy [50]. Schreiber *et al.* [51], in their retrospective cohort study on 2370 nulliparous women who underwent a vacuum-assisted delivery, reported that there were no significant differences in the rates of severe perineal lacerations between the episiotomy and the non-episiotomy group. A meta-analysis of fifteen studies regarding the effect of mediolateral episiotomy during vacuum-assisted delivery did not find a protective effect of episiotomy in nulliparous women (OR 0.68, [95% CI, 0.43–1.07]), whereas in multiparous women, it might even increase maternal morbidity (OR 1.27, [95% CI, 1.05–1.53]) [52].

In the literature, there is only one pilot RCT on routine versus selective use of episiotomy in operative vaginal deliv-

ery, including 200 nulliparous women. However, the study was underpowered to prove a significant change in the OASIS risk, mainly due to a high rate of episiotomy (52%) in the selective use group [53, 54]. Ankarcrona *et al.* [55] evaluated 63654 nulliparous women delivered with vacuum extraction. They showed that episiotomy could reduce the prevalence of OASIS from 15.5% to 11.8%. However, a high number of episiotomies had to be done to prevent one case of fourth-degree perineal injury due to its relative infrequency.

#### 4. Discussion

The prevention of perineal lesions during vaginal delivery is a priority for obstetricians. Some established risk factors may be identified before and during labor. Antenatal risk factors, such as ethnicity, primiparity, even though non-modifiable, should be identified and discussed with the woman.

Concerning modifiable intrapartum risk factors for OASIS, we identified labor augmentation, epidural anesthesia, and operative vaginal delivery. These factors can be considered modifiable, being not intrinsically present during labor and delivery but introduced by medical decision (labor augmentation and operative vaginal) or women preference (epidural anesthesia). Regarding labor augmentation and operative vaginal delivery, clinical audits aimed to improve obstetric practice reducing their inappropriate adoption may reduce the OASIS risk. However, when these interventions during labor cannot be avoided to guarantee fetal and maternal wellbeing, they could be referred to as non-modifiable factors. Regarding epidural analgesia, the increased risk of OASIS may be included in the patient counseling.

Since spontaneous vaginal births and operative vaginal delivery are two non-comparable conditions, we separately described the role of episiotomy in preventing severe perineal tears in these two types of delivery. In spontaneous vaginal births, routine use of episiotomy is not associated with relevant advantages compared with a selective one. Furthermore, the literature does not report absolute indications for episiotomy in spontaneous vaginal births. Similarly, the role of mediolateral episiotomy in preventing OASIS during an instrumental delivery has not been established due to the absence of properly designed and adequately sized RCT. Furthermore, performing a RCT is challenging, given the ethical difficulties (women will undergo an iatrogenic perineal injury without the certainty of avoiding OASIS) and feasibility of recruitment.

Although data from the literature show that episiotomy can reduce the prevalence of severe lacerations in assisted deliveries, the number of procedures needed to prevent one OASIS case is high [55]. Moreover, the group of patients who could benefit from episiotomy has not been identified yet. In this regard, evidence is further limited for specific subgroups of patients, such as in the case of female genital mutilations [56].

#### 5. Conclusions

Given that the “informed choice” is becoming increasingly relevant in modern obstetrics and that risk factors for OASIS are well known, women with high-risk factors for third- and fourth-degree lacerations should be identified and informed of their individual risk of complications.

#### Author contributions

All the authors contributed to the intellectual content of the study and approved the final version of the article. MF, RR, and MB—Study Conceptualization. SG, PCZ, and SU—Methodology. PCZ and IP—Resources and Data Curation. PCZ, SG, SL, and IP—Writing the Original Draft. PCZ, SG, IP, MB, MF, RR, SL, RDP, PP, FL, and SU—Writing, Review, and Editing. MF and SU—Visualization and Supervision. All authors contributed to the interpretation of results, as well as reviewed and approved the final version.

#### Ethics approval and consent to participate

Not applicable.

#### Acknowledgment

Not applicable.

#### Funding

This research received no external funding.

#### Conflict of interest

The author declares no conflict of interest. SG is the Editor Board Member of this journal, given his role as Editor Board Member, had no involvement in the peer-review of this article and has no access to information regarding its peer-review.

#### References

- [1] Waldman R. ACOG Practice Bulletin no. 198: Prevention and Management of Obstetric Lacerations at Vaginal Delivery. *Obstetrics & Gynecology*. 2019; 133: 185–185.
- [2] Dietz HP, Pardey J, Murray H. Pelvic floor and anal sphincter trauma should be key performance indicators of maternity services. *International Urogynecology Journal*. 2015; 26: 29–32.
- [3] Friedman AM, Ananth CV, Prendergast E, D'Alton ME, Wright JD. Evaluation of third-Degree and Fourth-Degree Laceration Rates as Quality Indicators. *Obstetrics & Gynecology*. 2015; 125: 927–937.
- [4] Oberwalder M, Connor J, Wexner SD. Meta-analysis to determine the incidence of obstetric anal sphincter damage. *British Journal of Surgery*. 2003; 90: 1333–1337.
- [5] Liebling RE, Swingler R, Patel RR, Verity L, Soothill PW, Murphy DJ. Pelvic floor morbidity up to one year after difficult instrumental delivery and cesarean section in the second stage of labor: A cohort study. *American Journal of Obstetrics & Gynecology*. 2004; 191: 4–10.
- [6] Fenner DE, Genberg B, Brahma P, Marek L, DeLancey JOL. Fecal and urinary incontinence after vaginal delivery with anal sphincter disruption in an obstetrics unit in the United States. *American Journal of Obstetrics & Gynecology*. 2003; 189: 1543–1549.
- [7] Bahl R, Strachan B, Murphy DJ. Pelvic floor morbidity at 3 years after instrumental delivery and cesarean delivery in the second

- stage of labor and the impact of a subsequent delivery. *American Journal of Obstetrics & Gynecology*. 2005; 192: 789–794.
- [8] Eason E, Labrecque M, Marcoux S, Mondor M. Anal incontinence after childbirth. *Canadian Medical Association Journal*. 2002; 166: 326–330.
- [9] Asif S, Mulic-Lutvica A, Axfors C, Eckerdal P, Iliadis SI, Fransson E, *et al*. Severe obstetric lacerations associated with postpartum depression among women with low resilience - a Swedish birth cohort study. *British Journal of Obstetrics and Gynaecology*. 2020; 127: 1382–90.
- [10] Scambia G, Raffaelli R, Baggio S, Garzon S, Laganà AS, Sartori E, *et al*. Needlestick injuries among obstetrics and gynecology trainees: A study design to investigate an underestimated priority. *Italian Journal of Gynaecology and Obstetrics*. 2018; 30: 27–36.
- [11] Franchi M, Garzon S, Laganà AS, Baggio S, Cromi A, Ghezzi F, *et al*. Needlestick injuries among obstetrics and gynecology trainees: A survey study. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2021; 259: 67–74.
- [12] Pergialiotis V, Bellos I, Fanaki M, Vrachnis N, Doumouchtsis SK. Risk factors for severe perineal trauma during childbirth: An updated meta-analysis. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2020; 247: 94–100.
- [13] Smith LA, Price N, Simonite V, Burns EE. Incidence of and risk factors for perineal trauma: a prospective observational study. *BMC Pregnancy and Childbirth*. 2013; 13: 59.
- [14] Goldberg J, Hyslop T, Tolosa JE, Sultana C. Racial differences in severe perineal lacerations after vaginal delivery. *American Journal of Obstetrics and Gynecology*. 2003; 188: 1063–1067.
- [15] Combs CA, Robertson PA, Laros RK. Risk factors for third-degree and fourth-degree perineal lacerations in forceps and vacuum deliveries. *American Journal of Obstetrics and Gynecology*. 1990; 163: 100–104.
- [16] King JR, Korst LM, Miller DA, Ouzounian JG. Increased composite maternal and neonatal morbidity associated with ultrasonographically suspected fetal macrosomia. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2012; 25: 1953–1959.
- [17] Raio L, Ghezzi F, Di Naro E, Buttarelli M, Franchi M, Dürig P, *et al*. Perinatal outcome of fetuses with a birth weight greater than 4500 g: an analysis of 3356 cases. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2003; 109: 160–165.
- [18] Stotland NE, Caughey AB, Breed EM, Escobar GJ. Risk factors and obstetric complications associated with macrosomia. *The International Journal of Gynecology & Obstetrics*. 2004; 87: 220–226.
- [19] Handa VL, Danielsen BH, Gilbert WM. Obstetric anal sphincter lacerations. *Obstetrics & Gynecology*. 2001; 98: 225–230.
- [20] Weissmann-Brenner A, Simchen MJ, Zilberberg E, Kalter A, Weisz B, Achiron R, *et al*. Maternal and neonatal outcomes of macrosomic pregnancies. *Medical Science Monitor*. 2012; 18: PH77–PH81.
- [21] Siggelkow W, Boehm D, Skala C, Grosslercher M, Schmidt M, Koelbl H. The influence of macrosomia on the duration of labor, the mode of delivery and intrapartum complications. *Archives of Gynecology and Obstetrics*. 2008; 278: 547–553.
- [22] Mazouni C, Menard J-P, Porcu G, Cohen-Solal E, Heckenroth H, Gamberre M, *et al*. Maternal morbidity associated with obstetrical maneuvers in shoulder dystocia. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2006; 129: 15–18.
- [23] Gauthaman N, Walters S, Tribe I-A, Goldsmith L, Doumouchtsis SK. Shoulder dystocia and associated manoeuvres as risk factors for perineal trauma. *International Urogynecology Journal*. 2016; 27: 571–577.
- [24] Rouse DJ, Weiner SJ, Bloom SL, Varner MW, Spong CY, Ramin SM, *et al*. Second-stage labor duration in nulliparous women: relationship to maternal and perinatal outcomes. *American Journal of Obstetrics and Gynecology*. 2009; 201: 357.e1–e7.
- [25] Janni W, Schiessl B, Peschers U, Huber S, Strobl B, Hantschmann P, *et al*. The prognostic impact of a prolonged second stage of labor on maternal and fetal outcome. *Acta Obstetrica Et Gynecologica Scandinavica*. 2002; 81: 214–221.
- [26] Cheng YW, Hopkins LM, Laros RK, Caughey AB. Duration of the second stage of labor in multiparous women: maternal and neonatal outcomes. *American Journal of Obstetrics and Gynecology*. 2007; 196: 585.e1–e6.
- [27] Ramm O, Woo VG, Hung Y, Chen H, Ritterman Weintraub ML. Risk Factors for the Development of Obstetric Anal Sphincter Injuries in Modern Obstetric Practice. *Obstetrics and Gynecology*. 2018; 131: 290–296.
- [28] Riskin-Mashiah S, O'Brian Smith E, Wilkins IA. Risk Factors for Severe Perineal Tear: can we do Better? *American Journal of Perinatology*. 2002; 19: 225–234.
- [29] Robinson JN, Norwitz ER, Cohen AP, McElrath TF, Lieberman ES. Episiotomy, operative vaginal delivery, and significant perinatal trauma in nulliparous women. *American Journal of Obstetrics and Gynecology*. 1999; 181: 1180–1184.
- [30] Grobman WA, Rice MM, Reddy UM, Tita ATN, Silver RM, Mallett G, *et al*. Labor Induction versus Expectant Management in Low-Risk Nulliparous Women. *New England Journal of Medicine*. 2018; 379: 513–523.
- [31] El-Sayed YY, Rice MM, Grobman WA, Reddy UM, Tita ATN, Silver RM, *et al*. Elective Labor Induction at 39 Weeks of Gestation Compared With Expectant Management: Factors Associated With Adverse Outcomes in Low-Risk Nulliparous Women. *Obstetrics & Gynecology*. 2020; 136: 692–697.
- [32] Loewenberg-Weisband Y, Grisaru-Granovsky S, Ioscovich A, Samueloff A, Calderon-Margalit R. Epidural analgesia and severe perineal tears: a literature review and large cohort study. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2014; 27: 1864–1869.
- [33] Hirsch E, Elue R, Wagner A, Nelson K, Silver RK, Zhou Y, *et al*. Severe perineal laceration during operative vaginal delivery: the impact of occiput posterior position. *Journal of Perinatology*. 2014; 34: 898–900.
- [34] Ponkey SE, Cohen AP, Heffner LJ, Lieberman E. Persistent fetal occiput posterior position: obstetric outcomes. *Obstetrics & Gynecology*. 2003; 101: 915–920.
- [35] Simkin P. The fetal occiput posterior position: state of the science and a new perspective. *Birth*. 2010; 37: 61–71.
- [36] Carroli G, Mignini L. Episiotomy for vaginal birth. *Cochrane Database of Systematic Reviews*. 2009; CD000081.
- [37] Goh R, Goh D, Ellepola H. Perineal tears - A review. *Australian Journal of General Practice*. 2018; 47: 35–38.
- [38] Kalis V, Laine K, de Leeuw J, Ismail K, Tincello D. Classification of episiotomy: towards a standardisation of terminology. *BJOG: an International Journal of Obstetrics & Gynaecology*. 2012; 119: 522–526.
- [39] American Academy of Pediatrics, American College of Obstetricians and Gynecologists, editors. Guidelines for perinatal care. Eighth edition. Elk Grove Village, IL: Washington, DC: American Academy of Pediatrics; The American College of Obstetricians and Gynecologists. 2017.
- [40] Murphy DJ, Strachan BK, Bahl R, Royal College of Obstetricians and Gynaecologists. Assisted Vaginal Birth: Green-top Guideline No. 26. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2020; 127: e70–e112.
- [41] Franchi M, Parissone F, Lazzari C, Garzon S, Laganà AS, Raffaelli R, *et al*. Selective use of episiotomy: what is the impact on perineal trauma? Results from a retrospective cohort study. *Archives of Gynecology and Obstetrics*. 2020; 301: 427–435.
- [42] Franchi M, Parissone F, Lazzari C, Garzon S, Laganà AS, Raffaelli R, *et al*. Rome wasn't built in a day: step-by-step improvement of obstetric care in the labor ward. *Archives of Gynecology and Obstetrics*. 2020; 301: 1591–1592.
- [43] Jiang H, Qian X, Carroli G, Garner P. Selective versus routine use of episiotomy for vaginal birth. *Cochrane Database of Systematic Reviews*. 2017; 2: CD000081.
- [44] Simó González M, Porta Roda O, Perelló Capó J, Gich Saladich I, Calaf Alsina J. Mode of Vaginal Delivery: a Modifiable Intrapartum Risk Factor for Obstetric Anal Sphincter Injury. *Obstetrics and Gynecology International*. 2015; 2015: 1–5.

- [45] Segal D, Baumfeld Y, Yahav L, Yohay D, Geva Y, Press F, *et al.* Risk factors for obstetric anal sphincter injuries (OASIS) during vacuum extraction delivery in a university affiliated maternity hospital. *The Journal of Maternal-Fetal & Neonatal Medicine.* 2020; 33: 999–1003.
- [46] Levin G, Friedman T, Shai D, Alcalay M, Ram E, Meyer R. The association between vacuum-assisted vaginal delivery and anal sphincter injury. *International Urogynecology Journal.* 2021; 32: 1771–1777.
- [47] Lund NS, Persson LKG, Jangö H, Gommesen D, Westergaard HB. Episiotomy in vacuum-assisted delivery affects the risk of obstetric anal sphincter injury: a systematic review and meta-analysis. *European Journal of Obstetrics & Gynecology and Reproductive Biology.* 2016; 207: 193–199.
- [48] van Bavel J, Hukkelhoven CWPM, de Vries C, Papatsonis DNM, de Vogel J, Roovers JWR, *et al.* The effectiveness of mediolateral episiotomy in preventing obstetric anal sphincter injuries during operative vaginal delivery: a ten-year analysis of a national registry. *International Urogynecology Journal.* 2018; 29: 407–413.
- [49] de Vogel J, van der Leeuw-van Beek A, Gietelink D, Vujkovic M, de Leeuw JW, van Bavel J, *et al.* The effect of a mediolateral episiotomy during operative vaginal delivery on the risk of developing obstetric anal sphincter injuries. *American Journal of Obstetrics and Gynecology.* 2012; 206: 404.e1–404.e5.
- [50] Macleod M, Strachan B, Bahl R, Howarth L, Goyder K, Van de Venne M, *et al.* A prospective cohort study of maternal and neonatal morbidity in relation to use of episiotomy at operative vaginal delivery. *BJOG: an International Journal of Obstetrics & Gynaecology.* 2008; 115: 1688–1694.
- [51] Schreiber H, Mevorach N, Sharon-Weiner M, Farladansky-Gershnel S, Shechter Maor G, Biron-Shental T. The role of mediolateral episiotomy during vacuum-assisted vaginal delivery with soft cup devices. *Archives of Gynecology and Obstetrics.* 2021; 303: 885–890.
- [52] Sagi-Dain L, Sagi S. Morbidity associated with episiotomy in vacuum delivery: a systematic review and meta-analysis. *BJOG: An International Journal of Obstetrics & Gynaecology.* 2015; 122: 1073–1081.
- [53] Murphy D, Macleod M, Bahl R, Goyder K, Howarth L, Strachan B. A randomised controlled trial of routine versus restrictive use of episiotomy at operative vaginal delivery: a multicentre pilot study. *BJOG: an International Journal of Obstetrics & Gynaecology.* 2008; 115: 1695–1703.
- [54] Serati M, Salvatore S, Braga A. Letter to the editor: selective episiotomy versus no episiotomy for severe perineal trauma: a systematic review with meta-analysis. *International Urogynecology Journal.* 2020; 31: 1715–1716.
- [55] Ankarcrona V, Zhao H, Jacobsson B, Brismar Wendel S. Obstetric anal sphincter injury after episiotomy in vacuum extraction: an epidemiological study using an emulated randomised trial approach. *BJOG: An International Journal of Obstetrics & Gynaecology.* 2021; 128: 1663–1671.
- [56] Auricchio V, Garzon S, Pomini P, Laganà AS, Casarin J, Cromi A, *et al.* Clitoral reconstructive surgery after female genital mutilation: A systematic review. *Sexual & Reproductive Healthcare.* 2021; 29: 100619.