

Prognostic Potential of the Preoperative Controlling Nutritional Status (CONUT) Score in Predicting Survival of Patients with Cancer: A Systematic Review

Soraya Kheirouri¹ and Mohammad Alizadeh²

¹Department of Nutrition, Tabriz University of Medical Sciences, Tabriz, Iran; and ²Nutrition Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

ABSTRACT

The nutritional status of a patient has prognostic potency concerning short- and long-term outcomes, including survival, in many diseases. The controlling nutritional status (CONUT) score is a method for assessing nutritional status and predicting outcomes of several diseases. This study sought to systematically identify the prognostic role of preoperative CONUT score on posttreatment overall survival (OS), recurrence-free survival (RFS), and cancer-specific survival (CSS) in patients with cancer. The PubMed, SCOPUS, and Google Scholar databases and Google were searched for all dates until December 2019. Original articles investigating the association of preoperative CONUT score with survival in cancer patients who underwent surgery were included. Duplicate and irrelevant reports were screened out and the remaining articles assessed for quality and data extracted during critical analysis. Results of multivariate analysis were used to evaluate the prognostic competence of CONUT score in predicting survival. The search method identified an initial 181 articles, of which 32 were included in the final analysis. Lower OS, CSS, and RFS rates were reported by 100%, 100%, and 87.0% of the included studies, respectively, in cancer patients with high CONUT scores. A prognostic role of the CONUT score for prediction of OS, CSS, and RFS in cancer patients was shown by 91.7%, 90.9%, and 52.6% of the studies, respectively. The receiver operating characteristic curve area under the curve (AUC) value of the CONUT score for predicting OS, CSS, and RFS was at an acceptable level (>0.5) in all studies with available AUC values ($n = 19$). Sixty percent (12 of 20) of the studies reported that high CONUT score was significantly related to lower BMI. The findings promote confidence that a high preoperative CONUT score is associated with poor survival rate and is an independent prognostic factor of OS and CSS in patients with various types of cancer. Evaluation of the preoperative CONUT score might help clinicians in decision-making with respect to surgical implications. *Adv Nutr* 2021;12:234–250.

Keywords: biomarker, cancer, cancer-specific survival, controlling nutritional status, overall survival, recurrence-free survival, prognosis

Introduction

A biomarker is a biological feature that is measured to indicate the body's biological situation or condition, objectively, in relation to, e.g., normal or pathogenic processes, or pharmacologic reactions to a therapeutic approach (1). In

clinical practice, biomarkers are often used for early diagnosis or monitoring of a disease; predicting the risk or outcome of a disease; measuring the effectiveness or harmfulness of a treatment regimen or other aspects of health; or screening of patients (2).

Prognostic biomarkers are baseline evaluations of disease features or patients' characteristics that can predict the risk or outcome of the disease in patients regardless of therapy (3). In cancers, a prognostic biomarker could be an attribute of the disease (such as size of the tumor or stage or grade of the cancer) or an individual characteristic of the patient (such as age, sex, or weight loss) that may influence the outcome. Prognostic biomarkers are beneficial in predicting the outcomes, developing a treatment plan, and decision-making regarding surgical or chemotherapy implications (4).

Supported by Tabriz University of Medical Sciences.

Author disclosures: The authors report no conflicts of interest.

Supplemental Table 1 is available from the "Supplementary data" link in the online posting of the article and from the same link in the online table of contents at <https://academic.oup.com/advances>.

Address correspondence to SK (e-mail: kheirouris@tbzmed.ac.ir) or MA (e-mail: mdalizadeh@tbzmed.ac.ir).

Abbreviations used: CA19-9, carbohydrate antigen 19-9; CEA, carcinoembryonic antigen; CONUT, controlling nutritional status; CSS, cancer-specific survival; DFS, disease-free survival; GPS, Glasgow Prognostic Score; NLR, neutrophil-to-lymphocyte ratio; OS, overall survival; PFS, progression-free survival; PLR, platelet-to-lymphocyte ratio; PNI, Prognostic Nutritional Index; RFS, recurrence/relapse-free survival; ROC, receiver operating characteristic.

Malnutrition is prevalent in cancer patients, which can affect short- and long-term outcomes. Nutritional status has a prognostic capacity for posttreatment long-term outcomes, including disease progression and survival, in patients (5). The controlling nutritional status (CONUT) score is an efficient nutritional screening tool for assessing the nutritional status of patients and is useful for early detection of undernutrition in all hospital inpatients. This tool is computed from the 3 clinical parameters of serum albumin, total cholesterol concentration, and total peripheral lymphocyte counts (6).

Multiple investigations have recently examined the prognostic capacity of the CONUT score on posttreatment complications, clinicopathological factors, and long-term outcomes in several types of diseases such as heart failure, liver disease, hypertension, and cancer (7–10). However, to our knowledge, the prognostic value of the CONUT score with regards to long-term outcomes has not yet been systematically reviewed in patients with cancer. The research question was “Is controlling nutritional status useful as a prognostic biomarker of survival in cancer patients who underwent surgery?” Therefore, this study aimed to systematically identify the prognostic role of preoperative CONUT score on posttreatment long-term outcomes including overall survival (OS), recurrence-free survival (RFS), and cancer-specific survival (CSS) in patients with cancer.

Methods

Search strategy and selection criteria

This review was conducted according to the guidelines indicated in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses protocols, 2015 statement. The PubMed, Google Scholar, and SCOPUS databases and Google were searched for all dates until December 2019. Journal articles and observational studies were searched. Original articles and congresses (if information of interest was available) in the English language that investigated the association of preoperative CONUT score with survival in cancer patients who underwent surgery were included. Inclusion criteria were as follows: articles that studied cancer patients who underwent surgery, and availability of postoperative survival information in the article. Research that studied cancer patients who underwent chemotherapy, radiotherapy, or immunotherapy without surgery, or with an unclear treatment method was excluded. Studies with postoperative CONUT score were also excluded. Studies that applied neoadjuvant therapy along with an operation were included in the present study if the effect of adjuvant therapy was considered in the multivariate analysis, otherwise they were excluded. Reviews, conferences, animal studies, editorials, and letters were also excluded. The following search terms were used: “controlling nutritional status” (in title) AND “survival” (in title/abstract) AND “cancer OR tumor OR carcinoma OR malignant” (in title/abstract). Reference lists of the articles were manually reviewed to identify further studies.

Screening of the articles

The extracted articles were saved in an EndNote file (Clarivate Analytics) and sorted to remove duplicate reports. The remaining titles and abstracts were reviewed to screen articles with the correct scope for the present review. The full texts of the screened articles were then critically analyzed separately for eligibility.

Quality assessment of the articles

The Newcastle-Ottawa scale for cohort studies was used for assessment of the quality of the included studies (11). Articles were categorized as good quality (if 3 or 4 stars in the selection domain AND 1 or 2 stars in the comparability domain AND 1 or 2 or 3 stars in the outcome/exposure domain were gained), fair quality (if 2 stars in the selection domain AND 1 or 2 stars in the comparability domain AND 2 or 3 stars in the outcome/exposure domain were achieved), or poor quality (if 0 or 1 star in the selection domain OR 0 stars in the comparability domain OR 0 or 1 star in the outcome/exposure domain were attained).

Data extraction

Full texts of the screened articles were carefully reviewed and data were extracted with regard to authors, year of publication, country, study design, type of cancer, type of treatment, number and age of patients, follow-up time, method and time of CONUT assessment, CONUT cutoffs, method of CONUT cutoff determination, efficacy of the CONUT score (sensitivity and specificity) for prognosis of survival, survival rates and their association with CONUT score, method of survival rate estimation, method of statistical analysis, confounding factors, and adjustments. Results of multivariate analysis were used to evaluate the prognostic effect of CONUT score on survival, in this review.

The time interval from surgery to radiological or histological detection of recurrence or metastasis or progression of disease was defined as RFS, disease-free-survival (DFS), or progression-free survival (PFS), which were all placed in 1 group (12, 13). The time interval from surgery to cancer-related death was defined as CSS (12, 14) and the time interval from surgery to overall death or last follow-up was defined as OS (12, 15). Studies that did not specify type of survival were considered as OS in this review.

Results

As **Figure 1** shows, initially 181 titles and abstracts were found using the search method (173 by databases, 5 by Google, and 3 by reference list searching). After elimination of duplicate studies, 62 articles remained. During the screening stage, 50 studies were found to be relevant to the study subject. During critical analysis, 18 articles were excluded because of not being written in English ($n = 3$), being conference papers ($n = 2$), being generic studies ($n = 2$), similarity ($n = 1$), low quality ($n = 1$), unclear treatment method ($n = 2$), postoperative CONUT score ($n = 1$), and applying adjuvant therapy without considering it in the multivariate analysis ($n = 6$). Eventually, 32 original studies




FIGURE 1 Flow diagram of the study. CONUT, controlling nutritional status.

(10, 12, 16–45) (W Yang, C Shou, J Yu, Q Zhang, X Liu, H Yu, X Lin, unpublished results, 2019) matched the study scope and, thus, were comprised in the final review and analysis (Figure 1).

Characteristics of the studies

As Table 1 shows, 62.5% ($n = 20$) of the included studies were performed in Japan, 31.3% ($n = 10$) in China, and 6.3% ($n = 2$) in South Korea. All the studies included were retrospective cohort investigations except the report by Huang et al. (22), a prospective cohort study. Of the articles, 96.9% (31 of 32) were published between 2017 and 2019.

In all included studies the CONUT score was evaluated based on serum albumin concentration, total lymphocyte count, and total cholesterol concentration in each patient. Of the studies, 90.6% (29 of 32) used the Kaplan–Meier method and log-rank test to estimate survival rates and 96.9% (31 of 32) used Cox proportional hazards regression models to calculate HRs and 95% CIs.

The receiver operating characteristic (ROC) curve and the AUC were used to evaluate the efficacy of the CONUT score for predicting survival (65.6%, 21 of 32) and to determine the optimal cutoff for the CONUT score. The highest Youden

index was used to attain the optimal cutoff of the CONUT score in 10 studies.

According to the Newcastle-Ottawa scale, all studies achieved good quality scores (Supplemental Table 1).

Relation between CONUT score and patients' survival

Twenty-four articles studied the OS rate in different CONUT groups, for which all the studies (100%) showed that patients with high CONUT scores had significantly shorter OS than those with low CONUT scores ($P < 0.05$). Twenty-four of the studies investigated the prognostic effect of high CONUT score with regards to OS. Based on the multivariate analysis, 91.7% (22 of 24) of the studies demonstrated that CONUT score was an independent prognostic factor of OS ($P < 0.05$).

Eleven studies investigated CSS, cancer-specific mortality (CSM), or disease-specific survival (DSS) rate in different CONUT groups and all the studies (100%) indicated that patients with a high CONUT score had significantly shorter CSS than those with a low CONUT score ($P < 0.05$). According to multivariate analysis, 90.9% (10 of 11) of the studies reported that CONUT score was an independent prognostic factor of CSS ($P < 0.05$).

Twenty-three studies investigated RFS, DFS, or PFS rate in different CONUT groups and 87.0% (20 of 23) of the

TABLE 1 Characteristics of the studies included and findings on prognostic potential (according to multivariate analysis) and efficacy of the CONUT score in predicting survival¹

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cut offs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Ahiko et al. (16)	Japan, retrospective cohort	Colorectal cancer (Stage I-IV), underwent surgery	830 (470 males, 360 females)	Median (range): 78 (75-94)	≥ 5 y	0-1 (n = 508) 2-3 (n = 249) ≥ 4 (n = 73)	Not indicated.	Not indicated.	*†‡	Higher CONUT group had the lowest 5-y OS rate ($P < 0.0001$). CONUT score was a significant prognostic factor of OS (≥ 4 vs. 0-1: HR: 2.24; 95% CI: 1.48, 3.30; $P < 0.001$) after adjustment for confounders.	BMI did not differ between CONUT groups.
Alkamine et al. (17)	Japan, retrospective	Lung adenocarcinoma, underwent surgery	109 (76 males, 33 females)	Mean (range): 72 (45-85)	4-13 y	High: ≥ 1 (n = 74) Low: 0 (n = 35)	ROC curve/—	AUC: 0.596 Sensitivity: 0.6711; specificity: 0.4375	*†‡	High-CONUT group had lower 5-y OS ($P = 0.04$) and DFS ($P = 0.01$) rates. CONUT score was independently associated with DFS (HR: 2.63; 95% CI: 1.33, 5.68; $P = 0.004$) and OS (HR: 2.64; 95% CI: 1.06, 7.80; $P = 0.04$).	High-CONUT group had lower BMI ($P = 0.025$).
Elghiaty et al. (12)	South Korea, retrospective cohort	Nonmetastatic clear cell renal cell carcinoma, underwent radical or partial nephrectomy	1046 (745 males and 301 females)	Median (range): 56 (46-64)	Median (range): 63 (43-87) mo	High: > 2 (n = 115) Low: ≤ 2 (n = 931)	ROC curve/Youden Index	AUC (based on OS) = 0.633 Sensitivity = 46.4%; specificity = 73.7% ($P = 0.001$)	*†‡	High-CONUT group had lower 3- and 5-y RFS ($P < 0.001$), CSS ($P = 0.006$), and OS ($P < 0.001$) rates. High CONUT score was an independent predictor of RFS (HR: 3.09; 95% CI: 1.45, 6.59; $P = 0.003$), CSS (HR: 4.66; 95% CI: 1.62, 13.39; $P = 0.004$), and OS (HR: 2.81; 95% CI: 1.44, 5.50; $P = 0.003$).	High-CONUT group had lower BMI ($P = 0.001$).

(Continued)

TABLE 1 (Continued)

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cut-offs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Harimoto et al. (18)	Japan, retrospective cohort	Hepatocellular carcinoma, underwent hepatic resection	2461 (1785 males, 676 females)	Mean ± SD: Low CONUT: 68.2 ± 10.1; High CONUT: 69.8 ± 9.2	Not indicated	High: ≥4 (n = 540) Low: ≤3 (n = 1921)	ROC curve/Youden index	AUC (based on OS): 0.580 Sensitivity: 31.3%; specificity: 91.6% (P < 0.01) AUC (based on CSS): 0.563 Sensitivity: 30.0%; specificity: 80.1% (P < 0.01) AUC (based on RFS): 0.536 Sensitivity: 63.5%; specificity: 41.9% (P < 0.01).	*†‡	High-CONUT group had lower OS and RFS rates (both, P < 0.01). Higher CONUT score was an independent predictor of poor OS (HR: 1.22; 95% CI: 1.06, 1.41; P = 0.006) and RFS (HR: 1.22; 95% CI: 1.06, 1.40; P = 0.006).	High-CONUT group had low BMI (P < 0.01).
Harimoto et al. (19)	Japan, retrospective	Hepatocellular carcinoma, underwent hepatic resection	357 (270 males, 87 females)	Mean ± SD: Low CONUT: 67.3 ± 10.7; High CONUT: 69.8 ± 8.5	≥5 y	High: >3 (n = 69) Low: ≤3 (n = 288)	ROC curve/Youden index	AUC (based on OS): 0.621 Sensitivity: 56.06%; specificity: 66.56% AUC (based on DSS): 0.651 Sensitivity: 65.7%; specificity: 59.5%.	*†‡	High-CONUT group had lower 5-y OS, RFS, and DSS rates (P < 0.01). Higher CONUT score was associated with poor OS (HR: 2.16; 95% CI: 1.25, 3.72; P = 0.03), but not with RFS.	BMI did not differ between CONUT groups.
Hirahara et al. (20)	Japan, retrospective	Esophageal cancer, underwent curative thoracoscopic esophagectomy	148 (132 males, 16 females)	Mean ± SD: CONUT 0: 65.6 ± 8.2; CONUT 1: 67.5 ± 8.4; CONUT 2-3: 65.3 ± 8.9	Not indicated	Normal nutrition: 0–1 (n = 70) Mild malnutrition: 2–4 (n = 62) Moderate–severe malnutrition: 5–12 (n = 16)	Not indicated.	Not indicated.	*†‡	CONUT was independently associated with worse prognosis for CSS (HR: 1.99; 95% CI: 1.07, 3.87), P = 0.03).	Not indicated.

(Continued)

TABLE 1 (Continued)

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cut offs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Hirahara et al. (21)	Japan, retrospective cohort	Gastric cancer, underwent curative gastrectomy	368 (254 males, 114 females)	Range: 36-91	Median: 35.3 mo	High: ≥ 3 (n = 105) Low: ≤ 2 (n = 263)	ROC curve/—	AUC (based on 5-y OS): 0.625 Sensitivity: 65.0%; specificity: 57.9%	* \diamond	High-CONUT group had lower 5-y OS rate (P < 0.001). Among all patients, OS was independently predicted by the CONUT score (HR: 2.25, P = 0.001). CONUT score was an independent prognostic factor for OS among the propensity score-matched subgroup (HR: 2.44; 95% CI: 1.46, 4.07; P < 0.001).	BMI did not differ between CONUT groups.
Huang et al. (22)	China, prospective cohort	Gastric cancer, underwent curative gastrectomy	357 (275 males, 82 females)	Mean \pm SD: 73.29 \pm 5.24	1 y	Normal: 0-1 (n = 153) Light: 2-4 (n = 168) Moderate and severe: ≥ 5 (n = 36)	Not indicated.	Not indicated.	\diamond	CONUT score was an independent predictor of postoperative 1-y survival (OR: 2.91; 95% CI: 0.91, 9.31; P = 0.02).	Moderate-severe CONUT group had lower BMI (P < 0.001).
Iseki et al. (23)	Japan, retrospective	Colorectal cancer (Stage II/III), underwent curative surgery	204 (112 males, 92 females)	Mean \pm SD: High CONUT: 66.09 \pm 9.23; Low CONUT: 71.13 \pm 11.57	8 y or until their deaths	High: ≥ 3 (n = 54) Low: ≤ 2 (n = 150)	ROC curve/—	AUC (based on 5-y CSS): 0.624 Sensitivity: 0.5263; specificity: 0.7622	* \diamond	High-CONUT group had lower 5-y CSS (P = 0.002) and RFS (P = 0.002) rates. CONUT score was an independent risk factor for CSS (OR: 4.21; 95% CI: 1.21, 13.35; P = 0.02), but not for RFS.	Not indicated.
Ishihara et al. (24)	Japan, retrospective cohort	Localized urothelial carcinoma treated with radical nephroureterectomy	107 (68 males, 39 females)	Mean \pm SD: Low CONUT: 72.7 \pm 9.98; High CONUT: 76.1 \pm 8.65	Mean \pm SD: 46.1 \pm 32.8; 25.5 \pm 18.4 mo	High: ≥ 3 (n = 24) Low: < 3 (n = 83)	ROC curve/Youden index	AUC (based on RFS): 0.588	* \diamond	High-CONUT group had lower 5-y RFS (P = 0.04), CSS (P = 0.004), and OS (P = 0.001) rates.	Not indicated.

(Continued)

TABLE 1 (Continued)

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cut offs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Kang et al. (25)	Korea, retrospective cohort	Renal cell carcinoma, underwent surgery	1881 (1361 males, 520 females)	Mean ± SD: Normal: 54.21 ± 12.17; Mild: 58.69 ± 12.80; Moderate to severe: 63.69 ± 12.83	Median (range): 41 (6–178) mo	High: ≥2 (n = 508) Low: 0–1 (n = 1373)	ROC curve/—	ROC curve	*†‡	CONUT score was an independent predictor of CSS (HR: 5.44; 95% CI: 1.95, 14.8; P = 0.002), OS (HR: 2.90; 95% CI: 1.18, 6.75; P = 0.02), and RFS (HR: 2.26; 95% CI: 0.97, 4.94; P = 0.058). High-CONUT group had shorter RFS (P = 0.02) and CSM (P < 0.001). High CONUT score was an independent predictor of CSM (HR: 1.89; 95% CI: 1.12, 3.20; P = 0.02), but not for RFS.	High-CONUT group had lower BMI (P < 0.001).
Kato et al. (26)	Japan, retrospective	Pancreatic adenocarcinoma, underwent resection	344 (207 males, 137 females)	Mean ± SD: 64.8 ± 9.9	Median (range): 29.1 (0.6–178.5) mo	High: ≥4 (n = 79) Low: <4 (n = 265)	ROC curve/—	AUC (based on 2-y survival): 0.614 (95% CI: 0.56, 0.67). Sensitivity: 30.6%; specificity: 88.6%	*†‡	High-CONUT group showed lower OS (P = 0.002) but not RFS. High CONUT score was an independent prognostic risk factor for OS (HR: 1.64; 95% CI: 1.19, 2.26; P = 0.003).	Not indicated.
Kuroda et al. (27)	Japan, retrospective cohort	Gastric cancer, underwent curative resection	416 (267 males, 149 females)	Median (range): 67.2 (25–94)	Median (range): 61.2 (1–134) mo	High: ≥4 (n = 62) Low: ≤3 (n = 354)	ROC curve/—	AUC (based on OS): 0.715 (95% CI: 0.68, 0.75) AUC (based on RFS): 0.658 (95% CI: 0.62, 0.70) AUC (based on CSS): 0.662 (95% CI: 0.61, 0.71)	*†‡	High-CONUT group had lower 5-y OS (P < 0.001), RFS (P = 0.02), and CSS (P = 0.02) rates. CONUT was an independent prognostic factor for OS (HR: 2.72; 95% CI: 1.74, 4.25; P < 0.001), but not for RFS and CSS.	High-CONUT group had lower BMI (P = 0.02).

(Continued)

TABLE 1 (Continued)

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cut offs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Liang et al. (28)	China, retrospective cohort	Soft-tissue sarcomas, underwent surgical resection	658 (393 males, 265 females)	Median (range): 43 (5–85)	Median (range): 103 (61–147) mo	High: ≥ 2 (n = 223) Low: 0–1 (n = 435)	ROC curve/Youden index	ROC curve	* \diamond	High-CONUT group had lower 5-y OS (P < 0.001) and DFS (P < 0.001) rates. High CONUT was an independent predictor of OS (HR: 1.86; 95% CI: 1.47, 4.14; P < 0.001) and DFS (HR: 1.63; 95% CI: 1.26, 2.11; P < 0.001).	Not indicated.
Lin et al. (29)	China, retrospective cohort	Hepatocellular carcinoma, underwent curative hepatectomy	380 (333 males, 47 females)	Median (range): 50 (19–80)	Median: 48.5 mo	High: ≥ 2 (n = 187) Low: < 2 (n = 193)	ROC curve/—	AUC (based on OS): 0.618 (95% CI: 0.567, 0.667) Sensitivity: 66.3%; specificity: 56.5%	* \diamond	High-CONUT group had lower 5-y OS (P < 0.001) and RFS (P = 0.02) rates. High CONUT was an independent prognostic indicator of decreased OS (HR: 2.40; 95% CI: 1.74, 4.25; P = 0.001), but not decreased RFS (HR: 1.36; 95% CI: 1.00, 1.85; P = 0.05).	Not indicated.
Miyata et al. (30)	Japan, retrospective	Intrahepatic cholangio-carcinoma, underwent curative hepatectomy	71 (45 males, 26 females)	Mean \pm SD: Low CONUT: 64.8 \pm 1.7; High CONUT: 69.1 \pm 1.9	Mean: 36.9 mo	High: ≥ 2 (n = 31) Low: < 2 (n = 40)	Not indicated.	Not indicated.	* \diamond	High-CONUT group had lower 1-, 3-, and 5-y OS (P = 0.01), but not RFS. High CONUT was an independent prognostic factor for OS (HR: 3.02; 95% CI: 1.4, 6.8; P = 0.007), but not for RFS.	High-CONUT group had lower BMI (P = 0.009).
Ryo et al. (31)	Japan, retrospective cohort	Gastric cancer, underwent gastrectomy	626 (435 males, 191 females)	Mean \pm SD: 67.9 \pm 10.9	Median: 49.2 mo or until death	High: ≥ 2 (n = 289) Low: < 2 (n = 337)	ROC curve/—	AUC (based on DFS): 0.656 Sensitivity: 0.66; specificity: 0.58	* \diamond	High-CONUT group had shorter OS (P < 0.0001) and DFS (P = 0.06) times. CONUT score was an independent prognostic factor for OS (HR: 1.74; 95% CI: 1.26, 2.41; P = 0.0007).	High-CONUT group had lower BMI (P < 0.0001).

(Continued)

TABLE 1 (Continued)

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cut offs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Shoji et al. (32)	Japan, retrospective	Non-small cell lung cancer, underwent surgery	138 (79 males, 59 females)	Mean (range): 68 (37–86)	Median (range): 58 (0–94) mo	High: ≥1 (n = 79) Low: 0 (n = 59)	ROC curve/—	AUC (based on CSS): 0.703 Sensitivity = 91.67%; specificity = 46.07%	*†‡	High-CONUT group had lower 5-y RFS (P = 0.046), CSS, (P = 0.01), and OS (P = 0.01) rates. CONUT score was an independent prognostic factor for CSS (RR: 6.06; 95% CI: 1.07, 113.94; P = 0.04).	Not indicated.
Song et al. (33)	China, retrospective cohort	Nonmetastatic renal cell carcinoma, underwent surgery	325 (231 males, 94 females)	Median (IQR): 57 (47–66)	Median (IQR): 64 (56.5–69) mo	High: ≥3 (n = 70) Low: <3 (n = 255)	ROC curve/Youden index	AUC (based on 5-y OS): 0.723, (P < 0.001) Sensitivity: 51.28%; specificity: 82.52%	*†‡	High-CONUT group had lower 5-y OS (P < 0.001), CSS (P < 0.001), and DFS (P < 0.001) rates. High CONUT was an independent risk factor for OS (HR: 3.36; 95% CI: 1.73, 6.56; P < 0.001), CSS (HR: 3.34; 95% CI: 1.59, 6.98; P = 0.001), and DFS (HR: 1.85; 95% CI: 1.07, 3.21; P = 0.03).	Not indicated.
Suzuki et al. (34)	Japan, retrospective	Gastric cancer, underwent curative resection	211 (141 males, 70 females)	≥75	Median (range): 47 (5–185) mo	Normal nutrition: (n = 75) Light malnutrition: (n = 100) Moderate or severe malnutrition: (n = 36)	Not indicated.	Not indicated.	*†‡	Higher-CONUT group had shorter OS (P < 0.001) and CSS (P < 0.001). CONUT score was an independent prognostic factor for OS (HR: 2.12; 95% CI: 1.18, 3.69; P = 0.01) and CSS (HR: 3.75; 95% CI: 1.30, 10.43; P = 0.01).	Higher CONUT group had lower BMI (P = 0.008).

(Continued)

TABLE 1 (Continued)

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cut offs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Takagi et al. (35)	Japan, retrospective cohort	Hepatocellular carcinoma, underwent hepatectomy	295 (241 males, 54 females)	Mean ± SD: 65.8 ± 10.4	Mean: 42.3 mo	High: ≥3 (n = 118) Low: ≤2 (n = 177)	ROC curve/—	AUC = 0.59	*†‡	High-CONUT group had lower 5-y RFS (P = 0.01) and OS (P = 0.006) rates. The CONUT score was an independent predictor of RFS (HR: 1.64; 95% CI: 1.15, 2.30; P = 0.006) and OS (HR: 2.50; 95% CI: 1.47, 4.23; P = 0.001).	BMI did not significantly differ between CONUT groups.
Takagi et al. (36)	Japan, retrospective cohort	Hepatocellular carcinoma, underwent hepatectomy	331 (269 males, 62 females)	Median (range): 67 (60–74)	1 mo	High: ≥5 (n = 30) Low: ≤4 (n = 301)	Not indicated.	Not indicated.	†‡	High-CONUT group had higher incidence of 30-d mortality (P < 0.001). High CONUT score was an independent predictor of in-hospital mortality after hepatectomy (HR: 9.41; 95% CI: 1.15, 77.4; P = 0.04).	BMI did not differ between CONUT groups (P > 0.05).
Takamori et al. (37)	Japan, retrospective	Malignant pleural mesothelioma	83 (66 males, 17 females)	Median (range): 59 (31–81)	—	High: ≥3 (n = 31) Low: ≤2 (n = 52)	ROC curve/—	AUC (based on 1-y survival): 0.772 Sensitivity: 73.1%; specificity: 64.0%	*†‡	High-CONUT group had lower OS and DFS rates (both, P < 0.001). High CONUT score was an independent predictive factor for OS (HR: 1.92; 95% CI: 1.17, 3.11; P = 0.01) and DFS (HR: 1.88; 95% CI: 1.14, 3.06; P = 0.01). High CONUT score was a prognostic factor of OS in patients who underwent surgery (HR: 4.86; 95% CI: 1.16, 19.14; P = 0.03).	Not indicated.

(Continued)

TABLE 1 (Continued)

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cut offs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Tokunaga et al. (38)	Japan, retrospective	Colorectal cancer, underwent curative resection	417 (247 males, 170 females)	Median (range): 68 (19–93)	Mean (range): 38.0 (1–115) mo	Normal: (n = 246) Light: (n = 127) Moderate: (n = 33) Severe: (n = 11)	Not indicated.	Not indicated.	*†‡	High-CONUT group (moderate/severe) had lower 5-y OS and RFS (both, P < 0.001). CONUT score was an independent prognostic factor for OS (moderate/severe vs. normal: HR: 5.92; 95% CI: 2.30, 14.92; P < 0.001; light vs. normal: HR: 2.74; 95% CI: 1.30, 5.87; P = 0.008), but not for RFS.	Moderate/severe CONUT group had lower BMI (P = 0.005).
Toyokawa et al. (39)	Japan, retrospective	Lung squamous cell carcinoma, underwent surgery	108 (96 males, 12 females)	Median (range): 71 (45–89)	≥ 5 y	High: ≥ 2 (n = 32) Low: 0–1 (n = 76)	ROC curve/—	AUC (based on OS): 0.590 Sensitivity: 0.786; specificity: 0.385	*†‡	High-CONUT group had lower 5-y DFS (P = 0.02) and OS (P = 0.006) rates. High CONUT score was an independent prognostic factor for DFS (HR: 1.90; 95% CI: 1.04, 3.37; (HR: 1.91; 95% CI: 0.92, 3.86; P = 0.08).	BMI did not differ between CONUT groups (P > 0.05).
Wang et al. (40)	China, retrospective	Malignant peritoneal mesothelioma	125 (46 males, 79 females)	Mean ± SD: 61.2 ± 9.3	Median (range): 8 (0.6–53) mo	High: ≥ 3 (n = 81) Low: ≤ 2 (n = 44)	ROC curve/—	AUC (based on OS): 0.861 (P < 0.001) Sensitivity: 0.784; specificity: 0.821	†	CONUT score was an independent predictive factor for OS (RR: 1.26; 95% CI: 1.16, 1.38; P < 0.001).	BMI did not differ between CONUT groups (P > 0.05).
Yamamoto et al. (41)	Japan, retrospective	Colorectal cancer (stage I–IV) underwent surgery	522 (291 males, 231 females)	—	≥ 5 y	High: ≥ 3 (n = 158) Low: < 3 (n = 364)	ROC curve/—	AUC (based on OS): 0.627 (P < 0.0001)	*†‡	High-CONUT group had a lower 5-y OS rate (P < 0.0001).	Not indicated.

(Continued)

TABLE 1 (Continued)

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cut-offs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Yang C et al. (42)	China, retrospective	Colorectal cancer, underwent curative resection	160 (90 males, 70 females)	Mean \pm SD: 58.4 \pm 11.8	Median (range): 30 (6–42) mo	High: ≥ 3 (n = 74) Low: < 3 (n = 86)	ROC curve/Youden index	AUC (based on CSS): 0.759 (P < 0.001) Specificity: 0.821; sensitivity: 0.625	* ∇ \diamond	High CONUT score was correlated with poor RFS (P < 0.001) and CSS (P < 0.001). CONUT score was an independent prognostic factor for RFS (HR: 2.02; 95% CI: 1.19, 3.43; P = 0.01) and CSS (HR: 3.45; 95% CI: 1.68, 7.10; P = 0.001). Higher-CONUT group had a lower RFS rate (P = 0.001). CONUT score was an independent prognostic factor for RFS (CONUT ≥ 5 vs. CONUT = 0–1; HR: 2.83; 95% CI: 1.46, 5.50; P = 0.002). High-CONUT group had shorter PFS before surgery (P < 0.05). High CONUT score was an independent prognostic factor for PFS (HR: 3.97; 95% CI: 1.05, 11.43; P = 0.004).	Not indicated.
W Yang, C Shou, J Yu, Q Zhang, X Liu, H Yu, X Lin, unpublished results, 2019	—, retrospective	Gastrointestinal stromal tumors, underwent resection	455 (222 males, 233 females)	Median (range): 57 (20–80)	Median (range): 110 (7–232) mo	Normal: 0–1 (n = 219) Light undernutrition: 2–4 (n = 196) Moderate–severe undernutrition: ≥ 5 (n = 40)	Not indicated.	Not indicated.	* ∇ \diamond	Higher-CONUT group had a lower RFS rate (P = 0.001). CONUT score was an independent prognostic factor for RFS (CONUT ≥ 5 vs. CONUT = 0–1; HR: 2.83; 95% CI: 1.46, 5.50; P = 0.002). High-CONUT group had shorter PFS before surgery (P < 0.05). High CONUT score was an independent prognostic factor for PFS (HR: 3.97; 95% CI: 1.05, 11.43; P = 0.004).	Not indicated.
Zhang et al. (43)	China, retrospective cohort	Metastatic prostate cancer, underwent surgery	94 males	Median (range): 71 (53–84)	Median (range): 16.31 (4.6–55.10) mo	High: ≥ 3 (n = 42) Low: 0–2 (n = 52)	X-tile program	Using the X-tile software	* \diamond	High-CONUT group had shorter PFS before surgery (P < 0.05). High CONUT score was an independent prognostic factor for PFS (HR: 3.97; 95% CI: 1.05, 11.43; P = 0.004).	BMI did not differ between CONUT groups (P > 0.05).
Zheng Z-F et al. (44)	China, retrospective	Gastric cancer, underwent radical gastrectomy	532 (403 males, 129 females)	Mean \pm SD: 61.1 \pm 11.5	Median (range): 60 (2–76) mo	Normal nutrition: n = 291 Light malnutrition: n = 183 Moderate or severe malnutrition: n = 58	Not indicated.	Not indicated.	* ∇ \diamond	Higher-CONUT group had the lowest 5-y OS (P = 0.006) and RFS (P = 0.02) rates. CONUT score was not associated with 5-y OS and RFS.	Higher-CONUT group had low BMI (P = 0.01).

(Continued)

TABLE 1 (Continued)

Reference	Country, study design	Type of cancer	Patients, n	Age, y	Follow-up time	CONUT score cutoffs	Determination of optimal cutoff values for CONUT score/index	CONUT score efficacy for predicting survival	Calculation method of survival rate, HR, and CONUT score ²	Findings	Relation of CONUT score and BMI
Zheng Y et al. (45)	China, retrospective	Renal cell carcinoma, underwent nephrectomy	635 (400 males, 235 females)	Mean ± SD: 61.71 ± 12.51	Median (range): 48.4 (29.3–80.1) mo	High: ≥2 (n = 349) Low: <2 (n = 286)	X-tile program	X-tile program	*†‡	High-CONUT group had shorter OS and CSS (both, $P < 0.0001$). CONUT score was an independent risk predictor of OS (HR: 3.01; 95% CI: 1.52, 5.95; $P = 0.001$) and CSS (HR: 3.00; 95% CI: 1.29, 6.98; $P = 0.01$).	High-CONUT group had low BMI ($P < 0.001$).

¹CONUT, controlling nutritional status; CSS, cancer-specific survival; DFS, disease-free survival; OS, overall survival; PFS, progression-free survival; RFS, recurrence/relapse-free survival; ROC, receiver operating characteristic.

²*Survival rate; †HR, cCONUT score. The Kaplan–Meier method and log-rank test were used to estimate survival rates. The Cox proportional hazards regression model was used to calculate HRs and 95% CIs. CONUT score was calculated from serum albumin and total cholesterol concentrations and total peripheral lymphocyte counts.

studies indicated that patients with a high CONUT score had significantly shorter RFS than those with a low CONUT score ($P < 0.05$). Nineteen studies evaluated the correlation between CONUT score and RFS. According to multivariate analysis, 52.6% (10 of 19) of the studies reported that the CONUT score was an independent prognostic factor for RFS ($P < 0.05$).

Efficacy of the CONUT score in prognosis of survival

Fourteen studies reported AUC values of ROC curves for OS, of which the minimum and maximum values were 0.58 and 0.86, respectively, among the studies. Eleven studies reported the sensitivity and specificity of CONUT score cutoffs for detection of OS. The minimum and maximum values of sensitivity were 30.6% and 82.0%, respectively, and for specificity were 38.5% and 91.6%, respectively, among the studies.

Four studies reported AUC values of ROC curves for RFS, ranging from 0.54 to 0.66 among the studies. Two studies reported CONUT score cutoffs' sensitivity (ranging from 63.5% to 66.0% among the studies) and specificity (ranging from 41.9% to 58.0% across the studies) for detection of RFS.

Six studies reported AUC values of ROC curves for CSS with minimum and maximum values of 0.56 and 0.76, respectively, across the studies. Five studies reported CONUT score cutoffs' sensitivity (ranging from 30.0% to 91.7% across the studies) and specificity (ranging from 46.1% to 80.1% across the studies) for detection of CSS.

Relation between BMI and CONUT score

Twenty studies measured the relation of BMI to CONUT score; 60.0% ($n = 12$) of the studies reported that the high-count group had significantly lower BMI ($P < 0.05$).

Discussion

Lower OS, CSS, and RFS rates in cancer patients with high CONUT scores were reported by 100%, 100%, and 86.7% of the studies, respectively. A prognostic role of the CONUT score for prediction of OS, CSS, and RFS in cancer patients was suggested by 91.7%, 90.9%, and 52.6% of the studies, respectively. The prognosis of outcomes (including tumor proliferation, patients' survival, and posttreatment complications) in patients with cancer has been demonstrated to be strongly correlated with aspects of the host's nutritional situation such as BMI, visceral obesity, and sarcopenia (46–49). The CONUT score is a nutritional assessment tool that is easily, cheaply, and objectively estimated from serum albumin, total cholesterol, and total lymphocyte count (6). A prognostic impact of the CONUT score on survival has also been reported in hospitalized elderly people (50) and in patients with heart failure (7), end-stage liver disease (8), hypertension (9), and peritoneal dialysis (51).

The AUC value of the ROC curve for the CONUT score was at an acceptable level (>0.5) for predicting OS, CSS, and RFS in all of the studies where AUC values were presented (12, 18, 27). This indicates that the preoperative CONUT score is a reliable and independent prognostic

marker of survival in cancer patients. Although sensitivity and specificity of the CONUT score for the prediction of survival were at an acceptable level (>50%) in most of the studies, the difference across the studies was possibly due to the adoption of various CONUT score cutoffs. It appears that a cutoff ≥ 3 has acceptable sensitivity and specificity in predicting survival.

Several studies have shown that the AUC of the CONUT score, according to the ROC curve, was significantly higher than the AUC of each component of CONUT including serum albumin and total cholesterol concentrations and total lymphocyte count (23, 27, 31, 38, 45). Furthermore, numerous studies have reported that individual components of the CONUT score, against the CONUT score itself, were not independent predictors of survival in patients with various types of cancers (18, 19, 23, 45, 52). Taken together, it may be interpreted that the CONUT score is a more valuable factor than its individual components for predicting survival.

In addition, several studies have compared prognostic accuracy of the CONUT score in predicting survival with other nutritional prognostic factors such as the Prognostic Nutritional Index (PNI) (29, 33, 53, 54). The PNI is calculated from the serum albumin concentration and total peripheral lymphocyte count. The prognostic value of the PNI for predicting survival in cancer patients has been confirmed in numerous studies (55–59). However, various studies have shown that the CONUT score had a higher ROC curve AUC for the prediction of survival than the PNI (29, 33, 53, 54). Moreover, contrary to the CONUT score, many studies have shown that the PNI was not an independent predictor of survival in patients with cancer (30, 36, 60). Iseki et al. (23) reported that the PNI was not an independent predictor of CSS in patients with colorectal cancer, whereas CONUT score was. Takagi et al. (35) also reported that the PNI, against CONUT score, was not an independent predictor of survival in patients with hepatocellular carcinoma. Y Zheng et al. (45) also demonstrated that the PNI was not an independent predictor of survival in patients with renal cell carcinoma, whereas CONUT score was. Collectively, considering the results of all the aforementioned studies, it is concluded that the CONUT score has higher prognostic accuracy than, and is preferable to, the PNI in predicting survival in different types of cancer.

The platelet-to-lymphocyte ratio (PLR), neutrophil-to-lymphocyte ratio (NLR), and Glasgow Prognostic Score (GPS) are inflammatory factors that have recently been suggested as prognostic factors for predicting inflammation and survival in many diseases and/or cancers (61–65). Multiple studies have compared the prognostic accuracy of the CONUT score with the inflammatory markers in predicting survival. Li et al. (53) indicated that CONUT had a higher AUC (of ROC) for the prediction of survival than the NLR in patients with breast cancer. Leem et al. (54) reported that the CONUT had a higher ROC curve AUC than the GPS in patients with lung cancer. Lin et al. (29) showed that the CONUT score had a higher AUC value than the inflammatory parameters (NLR and PLR) in patients

with hepatocellular carcinoma. Song et al. (33) reported that the CONUT AUC score for 5-y OS was higher than that of the NLR in patients with renal cell carcinoma. Toyokawa et al. (52) showed that the AUC of CONUT for predicting 3-y OS was higher than the AUCs of the PLR, NLR, and GPS in patients with esophageal cancer. Moreover, many studies have shown that the PLR, NLR, and GPS, against CONUT score, were not independent predictors of survival in patients with cancer (20, 29, 43, 45, 52). Putting together all the findings, it is suggested that the CONUT score, in comparison with the inflammatory markers, has higher prognostic accuracy and is superior in predicting survival in different types of cancer.

Carcinoembryonic antigen (CEA) and carbohydrate antigen 19-9 (CA19-9) are 2 cancer markers with prognostic significance for prediction of survival in patients with cancer (66, 67). CEA is a protein commonly produced in very low concentrations in the blood of adults. The blood concentration of CEA may be raised in some types of cancer (68, 69). CA19-9 is a blood type antigen and it may be increased in patients with gastrointestinal cancers (69). None of the included studies compared the prognostic accuracy (AUC value of the ROC curve) of these markers with that of the CONUT score in predicting survival in cancer patients. However, several of the included studies showed that CEA and CA19-9, against CONUT score, were not independent predictors of survival in patients with cancer (10, 23, 26, 30, 32, 38, 42, 60). More investigations are needed to clarify the prognostic accuracy of the CONUT score in predicting survival compared with the CEA and CA19-9 markers in patients with cancer.

Conclusions

The findings of the present study promote confidence that a preoperative high CONUT score is associated with a poor survival rate and is an independent prognostic factor of OS and CSS after surgery in patients with cancer.

Application of the findings.

This study suggests that the CONUT score is an easy and inexpensive indicator which not only could be used for evaluating nutritional status, but can also be beneficial as a prognostic marker of patients' survival in various types of cancer. Evaluation of the CONUT score might help clinicians in decision-making with respect to surgical implications.

Strengths of the study.

All the studies had good quality and, except 1, were published in the last 3 y. Nearly all of the studies used the same method to estimate survival rate and to analyze the correlation between CONUT score and survival. Further, in all studies included, the roles of main confounding factors including age, gender, BMI, stage of disease, tumor site, tumor number, tumor grade, tumor size, lymph node metastasis, presence of high-grade disease, adjuvant chemotherapy, immunohistochemistry, and operative procedures were considered in the analysis.

Limitations of the study.

All included studies were from East Asia (Japan, China, and South Korea), which may restrict the utilization of the findings among other nations and ethnic groups. Most of the studies did not indicate the role of BMI as a nutritional factor in prediction of survival. Several of the studies did not indicate the value of the AUC of the CONUT score for prediction of survival. The optimal cutoff of the CONUT score varied across the different studies.

Suggestions for future research.

The optimal cutoff of the CONUT score must be homogenized before its utilization in clinical practice. More investigations are needed to clarify the prognostic accuracy of the CONUT score in predicting survival compared with CEA and CA19-9 markers in patients with cancer.

Acknowledgments

The authors' responsibilities were as follows—both of the authors: were involved in the searching process, selection of the articles, data extraction, manuscript writing, and read and approved the final manuscript.

References

1. Naylor S. Biomarkers: current perspectives and future prospects. *Expert Rev Mol Diagn* 2003;3(5):525–9.
2. Carlomagno N, Incollingo P, Tammaro V, Peluso G, Rupealta N, Chiacchio G, Sandoval Sotelo ML, Minieri G, Pisani A, Riccio E, et al. Diagnostic, predictive, prognostic, and therapeutic molecular biomarkers in third millennium: a breakthrough in gastric cancer. *Biomed Res Int* 2017;7869802.
3. Oldenhuis CN, Oosting SF, Gietema JA, de Vries EG. Prognostic versus predictive value of biomarkers in oncology. *Eur J Cancer* 2008;44(7):946–53.
4. Lu J, Chen Y, Liu Y, Ding J, Piao Z, Liu W. Clinical significance of prognostic score based on age, tumor size, and grade in gastric cancer after gastrectomy. *Cancer Manag Res* 2018;10:4279–86.
5. Mantzorou M, Koutelidakis A, Theocharis S, Giaginis C. Clinical value of nutritional status in cancer: what is its impact and how it affects disease progression and prognosis? *Nutr Cancer* 2017;69(8):1151–76.
6. Ignacio de Ulíbarri J, González-Madroño A, de Villar NG, González P, González B, Mancha A, Rodríguez F, Fernández G. CONUT: a tool for controlling nutritional status. First validation in a hospital population. *Nutr Hosp* 2005;20(1):38–45.
7. Formiga F, Chivite D, Corbella X. Utility of the Controlling Nutritional Status (CONUT) score in patients admitted due to acute heart failure. *Int J Cardiol* 2017;235:203.
8. Fukushima K, Ueno Y, Kawagishi N, Kondo Y, Inoue J, Kakazu E, Ninomiya M, Wakui Y, Saito N, Satomi S, et al. The nutritional index 'CONUT' is useful for predicting long-term prognosis of patients with end-stage liver diseases. *Tohoku J Exp Med* 2011;224(3):215–19.
9. Sun X, Luo L, Zhao X, Ye P. Controlling Nutritional Status (CONUT) score as a predictor of all-cause mortality in elderly hypertensive patients: a prospective follow-up study. *BMJ Open* 2017;7(9):e015649.
10. Daitoku N, Miyamoto Y, Tokunaga R, Sakamoto Y, Hiyoshi Y, Iwatsuki M, Baba Y, Iwagami S, Yoshida N, Baba H. Controlling nutritional status (CONUT) score is a prognostic marker in metastatic colorectal cancer patients receiving first-line chemotherapy. *Anticancer Res* 2018;38(8):4883–8.
11. Penson DF, Krishnaswami S, Jules A, Seroogy JC, McPheeters ML. Newcastle-Ottawa quality assessment form for cohort studies [Internet]. In: Evaluation and treatment of cryptorchidism. Rockville

(MD): Agency for Healthcare Research and Quality (US); 2012. [Cited 2020 Aug 19]. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK115843/bin/appe-fm3.pdf>.

12. Elghiyat A, Kim J, Jang WS, Park JS, Heo JE, Rha KH, Choi YD, Ham WS. Preoperative controlling nutritional status (CONUT) score as a novel immune-nutritional predictor of survival in non-metastatic clear cell renal cell carcinoma of ≤ 7 cm on preoperative imaging. *J Cancer Res Clin Oncol* 2019;145(4):957–65.
13. National Cancer Institute. Definition of relapse-free survival – NCI Dictionary of Cancer Terms [Internet]. Bethesda (MD): National Cancer Institute; [cited 2020 Aug 19]. Available from: <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/event-free-survival>.
14. National Cancer Institute. Definition of cause-specific survival – NCI Dictionary of Cancer Terms [Internet]. Bethesda (MD): National Cancer Institute; [cited 2020 Aug 19]. Available from: <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/cause-specific-survival>.
15. National Cancer Institute. Definition of overall survival rate – NCI Dictionary of Cancer Terms [Internet]. Bethesda (MD): National Cancer Institute; [cited 2020 Aug 19]. Available from: <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/overall-survival-rate>.
16. Ahiko Y, Shida D, Horie T, Tanabe T, Takamizawa Y, Sakamoto R, Moritani K, Tsukamoto S, Kanemitsu Y. Controlling nutritional status (CONUT) score as a preoperative risk assessment index for older patients with colorectal cancer. *BMC Cancer* 2019;19:946.
17. Akamine T, Toyokawa G, Matsubara T, Kozuma Y, Haratake N, Takamori S, Katsura M, Takada K, Shoji F, Okamoto T, et al. Significance of the preoperative CONUT score in predicting postoperative disease-free and overall survival in patients with lung adenocarcinoma with obstructive lung disease. *Anticancer Res* 2017;37(5):2735–42.
18. Harimoto N, Yoshizumi T, Inokuchi S, Itoh S, Adachi E, Ikeda Y, Uchiyama H, Utsunomiya T, Kajiyama K, Kimura K, et al. Prognostic significance of preoperative controlling nutritional status (CONUT) score in patients undergoing hepatic resection for hepatocellular carcinoma: a multi-institutional study. *Ann Surg Oncol* 2018;25(11):3316–23.
19. Harimoto N, Yoshizumi T, Sakata K, Nagatsu A, Motomura T, Itoh S, Harada N, Ikegami T, Uchiyama H, Soejima Y, et al. Prognostic significance of preoperative controlling nutritional status (CONUT) score in patients undergoing hepatic resection for hepatocellular carcinoma. *World J Surg* 2017;41:2805–12.
20. Hirahara N, Matsubara T, Hayashi H, Takai K, Nakada S, Tajima Y. Prognostic importance of controlling nutritional status in patients undergoing curative thoracoscopic esophagectomy for esophageal cancer. *Am J Ther* 2018;25(5):e524–32.
21. Hirahara N, Tajima Y, Fujii Y, Kaji S, Kawabata Y, Hyakudomi R, Yamamoto T, Taniura T. Controlling Nutritional Status (CONUT) as a prognostic immunonutritional biomarker for gastric cancer after curative gastrectomy: a propensity score-matched analysis. *Surg Endosc* 2019;33(12):4143–52.
22. Huang Y, Huang Y, Lu M, Sun W, Sun X, Chen X, Li L, Chandoo A, Li L. Controlling nutritional status (CONUT) score is a predictor of post-operative outcomes in elderly gastric cancer patients undergoing curative gastrectomy: a prospective study. *Cancer Manag Res* 2019;11:9793–800.
23. Iseki Y, Shibusaki M, Maeda K, Nagahara H, Ohtani H, Sugano K, Ikeya T, Muguruma K, Tanaka H, Toyokawa T, et al. Impact of the preoperative controlling nutritional status (CONUT) score on the survival after curative surgery for colorectal cancer. *PLoS One* 2015;10(7):e0132488.
24. Ishihara H, Kondo T, Yoshida K, Omae K, Takagi T, Iizuka J, Tanabe K. Preoperative controlling nutritional status (CONUT) score as a novel predictive biomarker of survival in patients with localized urothelial carcinoma of the upper urinary tract treated with radical nephroureterectomy. *Urol Oncol* 2017;35(9):539.e9–16.

25. Kang HW, Seo SP, Kim WT, Yun SJ, Lee S-C, Kim W-J, Hwang EC, Kang SH, Hong S-H, Chung J, et al. Prognostic impact of nutritional status assessed by the controlling nutritional status (CONUT) score in patients with surgically treated renal cell carcinoma. *Nutr Cancer* 2018;70(6):886–94.
26. Kato Y, Yamada S, Suenaga M, Takami H, Niwa Y, Hayashi M, Iwata N, Kanda M, Tanaka C, Nakayama G, et al. Impact of the controlling nutritional status score on the prognosis after curative resection of pancreatic ductal adenocarcinoma. *Pancreas* 2018;47(7):823–9.
27. Kuroda D, Sawayama H, Kurashige J, Iwatsuki M, Eto T, Tokunaga R, Kitano Y, Yamamura K, Ouchi M, Nakamura K, et al. Controlling nutritional status (CONUT) score is a prognostic marker for gastric cancer patients after curative resection. *Gastric Cancer* 2018;21(2):204–12.
28. Liang Y, Hou T, Que Y, Zhao B, Xiao W, Zhang X, Zhou Z. Elevated controlling nutritional status (CONUT) score is associated with poor long-term survival in patients with low-grade soft-tissue sarcomas treated with surgical resection. *Clin Orthop Relat Res* 2019;477(10):2287–95.
29. Lin Z-X, Ruan D-Y, Jia C-C, Wang T-T, Cheng J-T, Huang H-Q, Wu X-Y. Controlling nutritional status (CONUT) score-based nomogram to predict overall survival of patients with HBV-associated hepatocellular carcinoma after curative hepatectomy. *Clin Transl Oncol* 2020;22(3):370–80.
30. Miyata T, Yamashita YI, Higashi T, Taki K, Izumi D, Kosumi K, Tokunaga R, Nakagawa S, Okabe H, Imai K, et al. The prognostic impact of controlling nutritional status (CONUT) in intrahepatic cholangiocarcinoma following curative hepatectomy: a retrospective single institution study. *World J Surg* 2018;42(4):1085–91.
31. Ryo S, Kanda M, Ito S, Mochizuki Y, Teramoto H, Ishigure K, Murai T, Asada T, Ishiyama A, Matsushita H, et al. The controlling nutritional status score serves as a predictor of short- and long-term outcomes for patients with stage 2 or 3 gastric cancer: analysis of a multi-institutional data set. *Ann Surg Oncol* 2019;26(2):456–64.
32. Shoji F, Haratake N, Akamine T, Takamori S, Katsura M, Takada K, Toyokawa G, Okamoto T, Maehara Y. The preoperative controlling nutritional status score predicts survival after curative surgery in patients with pathological stage I non-small cell lung cancer. *Anticancer Res* 2017;37(2):741–7.
33. Song H, Xu B, Luo C, Zhang Z, Ma B, Jin J, Zhang Q. The prognostic value of preoperative controlling nutritional status score in non-metastatic renal cell carcinoma treated with surgery: a retrospective single-institution study. *Cancer Manag Res* 2019;11:7567–75.
34. Suzuki S, Kanaji S, Yamamoto M, Oshikiri T, Nakamura T, Kakeji Y. Controlling nutritional status (CONUT) score predicts outcomes of curative resection for gastric cancer in the elderly. *World J Surg* 2019;43(4):1076–84.
35. Takagi K, Yagi T, Umeda Y, Shinoura S, Yoshida R, Nobuoka D, Kuise T, Araki H, Fujiwara T. Preoperative controlling nutritional status (CONUT) score for assessment of prognosis following hepatectomy for hepatocellular carcinoma. *World J Surg* 2017;41(9):2353–60.
36. Takagi K, Umeda Y, Yoshida R, Nobuoka D, Kuise T, Fushimi T, Fujiwara T, Yagi T. Preoperative controlling nutritional status score predicts mortality after hepatectomy for hepatocellular carcinoma. *Dig Surg* 2019;36(3):226–32.
37. Takamori S, Toyokawa G, Taguchi K, Edagawa M, Shimamatsu S, Toyozawa R, Nosaki K, Seto T, Hirai F, Yamaguchi M, et al. The controlling nutritional status score is a significant independent predictor of poor prognosis in patients with malignant pleural mesothelioma. *Clin Lung Cancer* 2017;18(4):e303–13.
38. Tokunaga R, Sakamoto Y, Nakagawa S, Ohuchi M, Izumi D, Kosumi K, Taki K, Higashi T, Miyamoto Y, Yoshida N, et al. CONUT: a novel independent predictive score for colorectal cancer patients undergoing potentially curative resection. *Int J Colorectal Dis* 2019;32(1):99–106.
39. Toyokawa G, Kozuma Y, Matsubara T, Haratake N, Takamori S, Akamine T, Takada K, Katsura M, Shimokawa M, Shoji F, et al. Prognostic impact of controlling nutritional status score in resected lung squamous cell carcinoma. *J Thorac Dis* 2017;9(9):2942–51.
40. Wang Q-Q, Zheng G-Q, Yang D-L, Liang Y-F, Yin W-J, Su S-S. Pretreatment controlling nutritional status score and lactate dehydrogenase as predictive markers of survival in patients with malignant peritoneal mesothelioma. *Nutr Cancer* 2018;70(8):1264–74.
41. Yamamoto M, Saito H, Uejima C, Taniao A, Tada Y, Matsunaga T, Sakamoto T, Honjo S, Ashida K, Fujiwara Y. Prognostic value of combined tumor marker and controlling nutritional status (CONUT) score in colorectal cancer patients. *Yonago Acta Med* 2019;62(1):124–30.
42. Yang C, Wei C, Wang S, Han S, Shi D, Zhang C, Lin X, Dou R, Xiong B. Combined features based on preoperative controlling nutritional status score and circulating tumour cell status predict prognosis for colorectal cancer patients treated with curative resection. *Int J Biol Sci* 2019;15(6):1325–35.
43. Zhang W, Wu Y, Zhang Z, Guo Y, Wang R, Wang L, Mao S, Zhang J, Yao X. Controlling nutritional status score: a new prognostic indicator for patients with oligometastatic prostate cancer. *Curr Probl Cancer* 2019;43(5):461–70.
44. Zheng Z-F, Lu J, Xie J-W, Wang J-B, Lin J-X, Chen Q-Y, Cao L-L, Lin M, Tu R-H, Zheng C-H, et al. Preoperative skeletal muscle index vs the controlling nutritional status score: which is a better objective predictor of long-term survival for gastric cancer patients after radical gastrectomy? *Cancer Med* 2018;7(8):3537–47.
45. Zheng Y, Bao L, Wang W, Wang Q, Pan Y, Gao X. Prognostic impact of the controlling nutritional status score following curative nephrectomy for patients with renal cell carcinoma. *Medicine (Baltimore)* 2018;97(49):e13409.
46. Renfro LA, Loupakis F, Adams RA, Seymour MT, Heinemann V, Schmoll HJ, Douillard JY, Hurwitz H, Fuchs CS, Diaz-Rubio E, et al. Body mass index is prognostic in metastatic colorectal cancer: pooled analysis of patients from first-line clinical trials in the ARCAD database. *J Clin Oncol* 2016;34(2):144–50.
47. Moon H-G, Ju Y-T, Jeong C-Y, Jung E-J, Lee Y-J, Hong S-C, Ha W-S, Park S-T, Choi S-K. Visceral obesity may affect oncologic outcome in patients with colorectal cancer. *Ann Surg Oncol* 2008;15(7):1918–22.
48. Akay S, Urkan M, Balyemez U, Erşen M, Taşar M. Is visceral obesity associated with colorectal cancer? The first volumetric study using all CT slices. *Diagn Interv Radiol* 2019;25(5):338–45.
49. Miyamoto Y, Baba Y, Sakamoto Y, Ohuchi M, Tokunaga R, Kurashige J, Hiyoshi Y, Iwagami S, Yoshida N, Watanabe M, et al. Negative impact of skeletal muscle loss after systemic chemotherapy in patients with unresectable colorectal cancer. *PLoS One* 2015;10(6):e0129742.
50. Cabré M, Ferreira C, Arus M, Roca M, Palomera E, Serra-Prat M. Evaluation of CONUT for clinical malnutrition detection and short-term prognostic assessment in hospitalized elderly people. *J Nutr Health Aging* 2015;19(7):729–33.
51. Zhou H, Chao W, Cui L, Li M, Zou Y, Yang M. Controlling Nutritional Status (CONUT) score as immune-nutritional predictor of outcomes in patients undergoing peritoneal dialysis. *Clin Nutr* 2020;39(8):2564–70.
52. Toyokawa T, Kubo N, Tamura T, Sakurai K, Amano R, Tanaka H, Muguruma K, Yashiro M, Hiraoka K, Ohira M. The pretreatment controlling nutritional status (CONUT) score is an independent prognostic factor in patients with resectable thoracic esophageal squamous cell carcinoma: results from a retrospective study. *BMC Cancer* 2016;16:722.
53. Li W, Deng Y, Ma G, Wu Q, Zhou Q. Controlling nutritional status (CONUT) score is a prognostic factor in patients with resected breast cancer. *Ann Oncol* 2019;30(Supplement_3):mdz095.062.
54. Leem AY, Jung JY, Lee SC, Kim EY, Lee SH, Chang J, Chung KY, Paik HC, Kim DJ, Lee JG. Prediction of postoperative pulmonary complications using preoperative controlling nutritional status (CONUT) score in patients with resectable non-small cell lung cancer. *Curr Dev Nutr* 2019;3(Suppl 1):nzz035.P12–032-19.
55. Ida N, Nakamura K, Saijo M, Kusumoto T, Masuyama H. Prognostic nutritional index as a predictor of survival in patients with recurrent cervical cancer. *Mol Clin Oncol* 2018;8(2):257–63.

56. Hsieh M-C, Rau K-M, Chiang P-H, Sung M-T, Lan J, Luo H-L, Huang C-C, Huang C-H, Su HY-L. Impact of prognostic nutritional index on overall survival for patients with metastatic urothelial carcinoma. *J Cancer* 2018;9(14):2466–71.
57. Sun J, Mei Y, Zhu Q, Shou C, Tjhoi WEH, Yang W, Yu H, Zhang Q, Liu X, Yu J. Relationship of prognostic nutritional index with prognosis of gastrointestinal stromal tumors. *J Cancer* 2019;10(12):2679–86.
58. Hirahara N, Tajima Y, Fujii Y, Kaji S, Yamamoto T, Hyakudomi R, Taniura T, Kawabata Y. Prognostic nutritional index as a predictor of survival in resectable gastric cancer patients with normal preoperative serum carcinoembryonic antigen levels: a propensity score matching analysis. *BMC Cancer* 2018;18(1):285.
59. Luvían-Morales J, González-Trejo S, Carrillo JF, Herrera-Goepfert R, Aiello-Crocifoglio V, Gallardo-Rincón D, Ochoa-Carrillo FJ, Oñate-Ocaña LF. Association of the prognostic nutritional index and overall survival in patients with colorectal cancer: a STROBE compliant retrospective cohort study. *Cancer Med* 2019;8(7):3379–88.
60. Liu X, Zhang D, Lin E, Chen Y, Li W, Chen Y, Sun X, Zhou Z. Preoperative controlling nutritional status (CONUT) score as a predictor of long-term outcome after curative resection followed by adjuvant chemotherapy in stage II-III gastric cancer. *BMC Cancer* 2018;18:699.
61. Lee YSG, Baradi A, Peverelle M, Sultani R, Adams H, Garlick J, Wilson AM. Usefulness of platelet-to-lymphocyte ratio to predict long-term all-cause mortality in patients at high risk of coronary artery disease who underwent coronary angiography. *Am J Cardiol* 2018;121(9):1021–6.
62. Wang J, Zhou X, He Y, Chen X, Liu N, Ding Z, Li J. Prognostic role of platelet to lymphocyte ratio in prostate cancer: a meta-analysis. *Medicine (Baltimore)* 2018;97(40):e12504.
63. Bojaxhiu B, Templeton AJ, Elicin O, Shelan M, Zaugg K, Walser M, Giger R, Aebersold DM, Dal Pra A. Relation of baseline neutrophil-to-lymphocyte ratio to survival and toxicity in head and neck cancer patients treated with (chemo-) radiation. *Radiat Oncol* 2018;13(1):216.
64. Yang L, Huang Y, Zhou L, Dai Y, Hu G. High pretreatment neutrophil-to-lymphocyte ratio as a predictor of poor survival prognosis in head and neck squamous cell carcinoma: systematic review and meta-analysis. *Head Neck* 2019;41(5):1525–35.
65. Jomrich G, Hollenstein M, John M, Baierl A, Paireder M, Kristo I, Ilhan-Mutlu A, Asari R, Preusser M, Schoppmann SF. The modified Glasgow prognostic score is an independent prognostic indicator in neoadjuvantly treated adenocarcinoma of the esophagogastric junction. *Oncotarget* 2018;9(6):6968–76.
66. Auclin E, Taieb J, Lepage C, Aparicio T, Faroux R, Mini E, Folprecht G, Salazar R, Benetkiewicz M, Banzi M, et al. Carcinoembryonic antigen levels and survival in stage III colon cancer: post hoc analysis of the MOSAIC and PETACC-8 trials. *Cancer Epidemiol Biomarkers Prev* 2019;28(7):1153–61.
67. Mirkin KA, Hollenbeak CS, Wong J. Prognostic impact of carbohydrate antigen 19-9 level at diagnosis in resected stage I–III pancreatic adenocarcinoma: a U.S. population study. *J Gastrointest Oncol* 2017;8(5):778–88.
68. Lee SU, Jwa E, Kim DY, Kim TH, Baek JY, Cha Y, Chang HJ, Oh JH. Analysis of unexplained carcinoembryonic antigen elevation after curative treatment of locally advanced rectal cancer. *Int J Clin Oncol* 2018;23(5):924–9.
69. Alkady MM, Abdel-Messeih PL, Nousseir NM. Assessment of serum levels of the adipocytokine chemerin in colorectal cancer patients. *J Med Biochem* 2018;37(3):313–9.