



Taibah University

Journal of Taibah University Medical Sciences

www.sciencedirect.com



Original Article

Isokinetic trunk muscle performance in adolescents with different body mass indices



Abdullah M. Al-Shenqiti, PhD^{a,*}, Hatem A. Emar, PhD^b, Fahad S. Algarni, PhD^c and Osama A. Khaled, PhD^d

^a Department of Physical Therapy, Faculty of Medical Rehabilitation Sciences, Taibah University, KSA

^b Department of Growth and Developmental Disorders in Children and its Surgery, Faculty of Physical Therapy, Cairo University, Egypt

^c Rehabilitation Science Department, College of Applied Medical Sciences, King Saud University, KSA

^d Basic Science Department, Faculty of Physical Therapy, Cairo University, Egypt

Received 22 November 2020; revised 6 March 2021; accepted 10 March 2021; Available online 24 April 2021

المخلص

أهداف البحث: تقييم ذروة عزم الدوران ومتوسط قوة العضلات القابضة والباسطة للعمود الفقري لدى المراهقين الذين تتراوح أعمارهم بين ١٢ و ١٨ عاماً ومعرفة ما إذا كان هناك اختلاف في أداء عضلات الجذع باختلاف نوع مؤشر كتلة الجسم.

طرق البحث: تم قياس ذروة عزم الدوران ومتوسط قوة العضلات القابضة والباسطة للعمود الفقري لدى ١٨٠ شخصاً (تتراوح أعمارهم من ١٢ إلى ١٨ عاماً) تم تصنيفهم إلى أربع مجموعات وفقاً لمؤشر كتلة الجسم. كما تم إجراء التقييم باستخدام جهاز الـاييزوكينيتيك حيث تم تقييم ذروة عزم الدوران ومتوسط قوة العضلات القابضة والباسطة للعمود الفقري بسرعات ٦٠ درجة و ١٢٠ درجة / ثانية. وتم تحليل البيانات باستخدام تحليل التباين متعدد الاتجاهات لتحديد ما إذا كانت هناك أي فروق ذات دلالة إحصائية بين المجموعات الأربع.

النتائج: أشارت نتائج تحليل التباين متعدد الاتجاهات إلى انخفاض في قوة عضلات الجذع مع زيادة مؤشر كتلة الجسم. وكذلك أوضحت النتائج ارتفاعاً ملحوظاً في عزم الدوران ومتوسط قوة عضلات الجذع في مجموعة الأشخاص ذوي الوزن الطبيعي مقارنة بالأشخاص الذين يعانون من نقص الوزن وزيادة الوزن والسمنة. وأيضاً أظهرت نتائج معامل ارتباط بيرسون وجود ارتباطاً معنوياً سلبياً بين الوزن وعزم دوران الذروة للعضلات القابضة والباسطة للعمود الفقري. وكان هناك ارتباطاً سلبياً ملحوظاً بين الوزن ومتوسط قوة هذه العضلات.

الاستنتاجات: خلصت هذه الدراسة إلى وجود علاقة سلبية بين مؤشر كتلة الجسم وعزم الدوران الأقصى ومتوسط قوة عضلات الجذع والعضلات القابضة

والباسطة للظهر. وبالتالي يجب أن تتضمن برامج إعادة التأهيل للمراهقين الذين يعانون من زيادة الوزن والسمنة برنامجاً لتقوية عضلات الظهر بهدف تحسين فاعلية تلك التدخلات العلاجية وبالتالي تحسين نوعية حياة هؤلاء الأشخاص.

الكلمات المفتاحية: التقييم باستخدام جهاز الـاييزوكينيتيك؛ قوة عضلات الظهر القابضة؛ قوة عضلات الظهر الباسطة؛ عزم الدوران الأقصى؛ مؤشر كتلة الجسم؛ المراهقين.

Abstract

Objective: This study aims to evaluate the isokinetic performance for the peak torque and average power of the spinal flexor and extensor muscles in adolescents aged 12 to 18 years. The study also analyses the differences between the trunk muscle peak torque and average power with body mass index (BMI).

Method: The peak torque and average power of the trunk flexor and extensor muscles were measured in 180 adolescents (aged 12–18 years). The participants were classified into four groups according to BMI. The Biodex isokinetic dynamometer in concentric mode at speeds of 60° and 120°/sec was used for assessment.

Results: One-way multivariate analysis of variance MANOVA results demonstrated a significant difference in trunk muscle peak torque and average power with different BMI ($F = 14.692, p = 0.0005$). A Pearson's correlation analysis demonstrated a significantly negative correlation between weight and peak torque of trunk flexors and extensors ($r = -0.43, p = 0.0001$; $r = -0.31, p = 0.007$, respectively). Finally, the results showed a negative correlation between weight and average power

* Corresponding address:

E-mail: Abdullahalshenqiti@gmail.com (A.M. Al-Shenqiti)

Peer review under responsibility of Taibah University.



Production and hosting by Elsevier

of trunk flexors and extensors ($r = -0.54, p = 0.0001$; $r = -0.31, p = 0.007$).

Conclusion: In this study, overweight and obese adolescents are found to be correlated with decreased trunk muscle torque and power. Thus, therapeutic interventions for overweight and obese adolescents, along with exercise training programmes, may help improve muscle performance including peak torque and power. Finally, these measures may enhance the quality of life of such adolescents.

Keywords: Adolescents; BMI; Isokinetic evaluation; Peak torque; Trunk flexors and extensors

© 2021 The Authors.

Production and hosting by Elsevier Ltd on behalf of Taibah University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Recently, increased body mass index (BMI) at young age has emerged as a prevalent health concern.¹ Additionally, in Saudi children and adolescents, overweight and obesity have demonstrated alarming pervasiveness of 13.4% and 18.2%, respectively.²

Being overweight or obese can have many negative effects on the well-being of children and adolescents and lead to various health problems, including diabetes, cancer, musculoskeletal problems and respiratory and cardiac issues.^{3,4} A recent study on 343 participants with different BMI found that the prevalence of low back pain among obese adolescents (age 12–15 years) was 67%.⁵

Researchers have examined the impact of obesity on maximum isotonic,⁶ isometric,⁷ and isokinetic^{8–10} strength in different age groups from adolescents to the elderly. They concluded that muscle strength is lower when normalised to total body mass. Additionally, relative muscle strength of knee extension, trunk flexion, and handgrip in relation to body mass was about 10% lower in obese persons.¹¹

Pajoutan et al.¹² conducted a study to investigate the effect of body composition on trunk extensor muscles' maximum capacity across all levels of BMI-defined categories and over a wider age range. They found that absolute back muscle strength was comparable between BMI categories, while relative strength to body mass in obese participants was about 38% lower than that in people with normal weight and approximately 26% less than the overweight group.

Early detection and treatment of childhood musculoskeletal abnormalities is a vital intervention that should be undertaken by healthcare professionals.^{13,14} Efficient prevention programmes for children and adolescents can be established based on early detection of musculoskeletal problems, including spinal conditions.¹³

Measuring the strength and performance of trunk muscles is a challenging issue.^{15,16} However, this examination is

crucial for research and clinical practice to attain certain goals, including providing guidance to rehabilitation and prevention programmes.^{17,18}

Many objective tools can measure and train the strength of back muscles. Isokinetic testing ranks top among such methods.^{15,16} The assessment of muscle strength of trunk musculature at various positions, velocities, and modes shows a high level of safety,¹⁹ reliability,²⁰ and sensitivity that is sufficient for the detection of muscle weakness²¹ and achieving the goals of rehabilitation.²² Thus, isokinetic dynamometry is an effective and reliable tool that objectively assesses the strength of specific muscle groups.²³ In a systematic review, Estrázulas et al.²⁴ concluded that the isokinetic testing of trunk muscles showed a high level of reliability; thus, they recommended the use of isokinetic measurement of trunk muscle performance in clinical settings.

Although muscle strength is considered an excellent indicator of general health when based on reliable measurements, most studies involving isokinetic measurements of trunk muscle performance recruit adult participants.^{25–31} However, few have measured trunk muscle strength isokinetically in normal healthy children and adolescents.³² To the best of our knowledge, there is a lack of published information on the values of isokinetic peak torque and the power of trunk muscles in Saudi children and adolescents and the relationship between BMI and these parameters.

Given the well-documented importance of the good performance of trunk muscles for optimal physical function, along with our limited knowledge of the effects of body weight on trunk muscle peak torque and power in young people, we conduct the present study to achieve the following objectives: (1) evaluate the isokinetic peak torque and mean power of trunk flexors and extensors in Saudi adolescents aged 12 to 18 years, (2) compare the isokinetic peak torque and mean power of trunk flexors and extensors in these adolescents with different BMI, and (3) determine the effects of weight on trunk muscle peak torque and power. We hope that our results provide strength-test administrators and teams working in rehabilitation with the normal values of peak torque and mean power of trunk flexors and extensors, as well as knowledge of variation in these parameters depending on weight. We hypothesise that trunk muscle peak torque and power is associated with body mass and that increasing mass reduces trunk muscle peak torque and power in adolescents.

Materials and Methods

Participants

The sample size of the present school-based cross-sectional study was 180 healthy male students who were recreationally active but non-participants in physical exercise. They were recruited from intermediate and secondary schools from Almadinah Almunawwarah city, KSA. The participants were selected using a multistage stratified sampling method. No participant reported practising sports or exercising for over half an hour. Adolescents were included in the study if they were registered in public schools, were in

the classroom on the day of measurements, and were aged 12 to 18 years. Participants with postural abnormalities recognised by observation, a history of musculoskeletal trauma to spine or hip areas, a history of surgery in the abdominal region and spine, lower back pain, trauma, or deformity of the knees were excluded from the study.

Procedures

Weight (kg) and height (m) were measured to the nearest 0.5 kg and 0.5 cm, respectively, using Detecto Physician's Scale Model 2392, USA. BMI was calculated by dividing weight in kilograms with height in square metres. Participants were considered obese if the BMI specific for their age and sex was at the 95th percentile or more of the growth charts of the Centres for Disease Control and Prevention. If the BMI was between the 85th and 95th percentiles, the participant was classified as overweight. If the BMI was between the 85th percentile and the 5th percentiles, the participant was classified as having normal weight. Participants were classified as underweight when their BMI was lower than the 5th percentile.³³

Testing protocol description

Before commencing with testing, a warm-up activity was performed, such as static cycling for 5 min at level two resistance.³⁴ Back flexor/extensor strength and power were measured on a Biodex isokinetic dynamometer 4 Pro (Biodex Corporation, Shirley, NY, USA). Participants were positioned on the back attachment designed to measure trunk extension–flexion. The spine was set upright and the hips and knees are positioned in 90° flexion position with the thighs aligned parallel to the ground. The mechanical axis of the dynamometer was aligned with the anatomical axis of the trunk, which is represented by a line joining the anterior superior iliac spines. This position is regarded as the anatomic reference position (Figure 1). Adjustable pads are positioned posterior to the head, the upper trunk, the pelvis, and on the anterior shaft of the tibia to fix the participant to the back attachment. The upper trunk, the pelvis, and the thighs are stabilised using straps. The range of movement of the back was set at 50°, which is composed of a flexion component of 30° (–30°) and an extension component of 20° (+20°), in relation to the previously mentioned anatomic reference angle (0°) (Figure 1). A trunk range of motion that does not exceed 50° helps in isolating the lumbar spine motion with decreasing flexion and extension of the hip joint.³⁵ Furthermore, the previously described alignment of mechanical and anatomical axes, the presence of the pad support behind the pelvis, and the stabilising belt on the pelvis limit unwanted motion at the hip joint during the measurements.

The measurement protocol began with the person in a flexed position, with their arms and hands crossed over the thorax, and included three sets of five maximum sequential concentric lumbar flexions and extensions with 60°/s and 120°/s angular velocities. In this study, each participant was

verbally encouraged to exert maximum effort throughout the whole protocol.

Data analysis

Statistical analysis was performed using SPSS 21.0 (Statistical Package for Social Sciences, IBM, New York). Normality of the data was verified using the Shapiro–Wilk test. Thus, parametric tests were applied to analyse quantitative data. One-way multiple analysis of variance (one-way MANOVA) with Tukey's post-hoc tests was used to observe differences in peak torque and average power of trunk flexors and extensors at 60 and 120 °/s in underweight, normal-weight, overweight, and obese conditions. Pearson's correlation was used to investigate the correlation between weight and peak torque and the average power of trunk flexors and extensors. Statistical significance was set at $p < 0.05$.

Results

Demographic data of the participants is presented in Table 1. Our sample includes 180 Saudi male participants (45 participants for each subgroup). The mean \pm SD age, weight, and height of the entire sample were 15.374 ± 2.015 years, 61.733 ± 27.560 kg, and 157.233 ± 16.921 cm, respectively.

The one-way MANOVA showed a statistically significant difference in peak torque and average power of trunk extensors and flexors at the two selected speeds based on the participants' BMI ($F(3, 176) = 14.692, p = 0.0005$; Wilks' Lambda = 0.450, partial $\eta^2 = 0.408$) (see Table 2).

Additionally, BMI was found to have a statistically significant effect on the peak torque of trunk extensors at 60 degrees/second ($F(3, 176) = 9.759; p < .0005$; partial $\eta^2 = 0.192$) and the peak torque of trunk flexors at 60 degrees/second ($F(3, 176) = 7.410; p = .0001$; partial $\eta^2 = 0.112$).

Furthermore, the peak torque of trunk extensors at 60 degrees/second for the underweight condition was significantly lower than that for the normal-weight condition (mean difference = 19.54). Moreover, the normal-weight condition is significantly higher than the overweight condition (mean difference = 29.931). Furthermore, the normal-weight condition significantly differs from the obese condition (mean difference = 32.373). Therefore, these results



Figure 1: flexion–extension trunk motion on the isokinetic machine through a 50° ROM.

Table 1: Demographic data of participants.

Variable	Mean \pm SD				F	p
	Underweight (N = 45)	Normal weight (N = 45)	Overweight (N = 45)	Obese (N = 45)		
Age (years)	15.184 \pm 2.606	15.768 \pm 2.029	15.079 \pm 2.261	15.755 \pm 2.165	1.166	0.3241
Weight (kg)	35.066 \pm 8.2204	53.311 \pm 16.641	70.488 \pm 8.04779	88.066 \pm 33.066	61.975	0.000
Height (cm)	146.822 \pm 10.149	148.511 \pm 8.381	147.177 \pm 8.7549	147.422 \pm 11.081	0.256	0.8570

Table 2: Descriptive statistics of peak torque and average power of trunk flexors and extensors based on BMI type.

Variable	Mean \pm SD				F	p
	Underweight (N = 45)	Normal weight (N = 45)	Overweight (N = 45)	Obese (N = 45)		
PTTE at 60°/s	89.573 \pm 23.726	109.113 \pm 28.161	79.188 \pm 32.444	76.740 \pm 37.135	10.312	0.000*
PTTF at 60°/s	82.055 \pm 24.884	136.286 \pm 40.248	78.793 \pm 26.8054	77.637 \pm 29.247	38.228	0.000*
APTE at 60°/s	36.104 \pm 21.032	46.222 \pm 19.363	35.6867 \pm 25.39903	32.942 \pm 19.001	3.352	0.020*
APTF at 60°/s	28.600 \pm 12.418	32.635 \pm 11.834	24.0622 \pm 19.64635	20.940 \pm 17.683	4.767	0.003*
PTTE at 120°/s	62.075 \pm 23.234	102.688 \pm 40.1513	66.404 \pm 36.565	53.280 \pm 23.090	21.165	0.000*
PTTF 120°/s	72.757 \pm 23.311	83.137 \pm 24.04366	66.6733 \pm 28.41589	59.9867 \pm 34.449	5.5824	0.001*
APTE at 120°/s	24.686 \pm 12.827	36.8933 \pm 8.998	28.5067 \pm 11.7839	27.49267 \pm 9.448	10.4903	0.000*
APTF at 120°/s	17.271 \pm 6.080	27.0133 \pm 9.392	19.4822 \pm 5.283	18.1822 \pm 4.790	11.221	0.000*

PTTE at 60°/s: Peak torque of trunk extensors at 60 degrees/second, PTTF at 60°/s: Peak torque of trunk flexors at 60 degrees/second, PTTE at 120°/s: Peak torque of trunk extensors at 120 degrees/second, APTF at 120°/s: Average power of trunk flexors at 120 degrees/second, APTF at 120°/s: Average power of trunk flexors at 120 degrees/second. APTE°/s at 120°/s: Average power of trunk extensors at 120 degrees/second.

* = Significant difference ($p < 0.05$).

suggest that overweight and obesity impact the peak torque of trunk extensors at 60 degrees/second was higher in normal weight condition. Specifically, our results suggest that the peak torque decreases with increase in body weight (Figure 2).

Moreover, the peak torque of trunk flexors at 60 degrees/second for the underweight condition differed significantly from that for the obese condition (mean difference = 4.418). Additionally, the normal-weight condition significantly differed from the overweight condition (mean difference = 57.493) and from the obese (mean difference = 58.649) in this regard. Thus, these results suggest that overweight and obesity influence the peak torque of trunk flexors at 60 degrees/second. Specifically, when body weight increases, the peak torque decreases.




Figure 2: Peak torque of trunk extensors at 60 degrees/second in four groups.

BMI also had a statistically significant effect on the average power of trunk extensors at 60 degrees/second ($F(3, 176) = 3.352$; $p = 0.02$; partial $\eta^2 = 0.37$) and average power of trunk flexors at 60 degrees/second ($F(2, 57) = 14.30$; $p < .0005$; partial $\eta^2 = .054$). Additionally, the results revealed that the average power of trunk extensors at 60 degrees/second for the normal-weight condition differed significantly from that for the obese condition (mean difference = 13.28) (Figure 3). Furthermore, the average power of trunk flexors at 60 degrees/second for the normal-weight condition differed significantly from that for the obese condition (mean difference = 11.695).

Similarly, a statistically significant effect of BMI on the peak torque of trunk extensors at 120 degrees/second ($F(3, 176) = 21.165$; $p = .000$; partial $\eta^2 = 0.265$) and peak torque of trunk flexors at 120 degrees/second ($F(3, 176) = 43.518$; $p = .000$; partial $\eta^2 = 0.426$) was observed. The peak torque of trunk extensors at 120 degrees/second for the normal-weight condition differed significantly from that for the underweight condition (mean difference = 40.613) Furthermore, normal weight differs significantly from overweight (mean difference = 36.284), and from the obese (mean difference = 49.408). Similarly, the peak torque of trunk flexors at 120 degrees/second was higher in normal-weight condition was significantly different from that for the overweight condition (mean difference = 16.464) and the obese condition (mean difference = 23.151) (Figure 4).

Additionally, our results showed that BMI had a statistically significant effect on the average power of trunk extensors at 120 degrees/second ($F(3, 156) = 6.851$; $p = 0.0001$; partial $\eta^2 = 0.31$) and average power of trunk flexors at 120 degrees/second ($F(3, 176) = 2.452$; $p = 0.038$; partial $\eta^2 = 0.057$). The average power of trunk extensors at 120




Figure 3: Average power of trunk extensors at 60 degrees/second.




Figure 4: Peak torque of trunk flexors at 120 degrees/second.

degrees/second for the underweight condition differed significantly from that for the normal-weight condition (mean difference = 12.207). The normal-weight condition differed significantly from the overweight condition (mean difference = 8.387) and from the obese condition (mean difference = 9.401) in this regard as well. The average power of trunk flexors at 120 degrees/second for the underweight condition differed significantly from that for the normal-weight condition (mean difference = 9.742). The normal-




Figure 5: Average power of trunk flexors at 120 degrees/second.

weight condition differed significantly from the overweight condition (mean difference = 7.531) and from the obese condition (mean difference = 8.831) (Figure 5) in this regard as well.

Correlation between weight and peak torque and average power of trunk flexors and extensors

Pearson's correlation analysis demonstrated a significant negative correlation between weight and the peak torque of trunk flexors and extensors at 60 degrees/second ($r = -0.43$, $p = 0.0001$; $r = -0.31$, $p = 0.007$, respectively). Additionally, there was a notable negative correlation between weight and average power of trunk flexors and extensors at 60 degrees/second ($r = -0.54$, $p = 0.0001$; $r = -0.31$, $p = 0.007$). Similarly, a significant negative correlation between weight and the peak torque of trunk flexors and extensors at 120 degrees/second ($r = -0.51$, $p = 0.0001$; $r = -0.53$, $p = 0.0001$, respectively). Furthermore, a significant negative correlation was found between weight and the average power of trunk flexors and extensors at 120 degrees/second ($r = -0.21$, $p = 0.04$; $r = -0.24$, $p = 0.007$).

Discussion

The current study aimed to compare the respective strength and average power of trunk flexors and extensors in adolescents aged 12–18 years with different BMI using an isokinetic dynamometer.

In general, our findings show that increased BMI has a significant negative relationship with the performance of spinal flexors and extensors, including peak torque and average power. Normal weight adolescents' values of muscle performance are higher than those of the overweight and obese adolescents, with significant differences.

In the present study, the isokinetic measurement of trunk flexors and extensors was performed in a seated position as this position seems to be minimally influenced by the contraction of the muscles of the hip, leading to reduction of stress on the lumbar region.³⁶ Therefore, this position is more suitable for the proper assessment of the strength of trunk flexors and extensors.

The results show that torque increases when moving towards lower velocities. This can be explained by the existence of an inverse relationship between muscle force and velocity of concentric muscle contraction. The slower the contraction of the muscle, the larger the tension it produces, and vice versa. Termed as the force–velocity relationship, this is considered as a basic principle of the physiology of skeletal muscle.³⁷

Our results are in line with those of Pajoutan et al.¹² who examined the effect of obesity on the muscle performance of back extensor musculatures. They recruited obese, overweight, and normalweight groups and had them undertake four sets of maximum volitional isometric contractions of back extensors, repeated thrice. They concluded that obesity negatively affects muscle performance in activities demanding stronger contractions. Generally, obesity at different ages results in reduced peak strength when expressed in relation to body mass.^{38,39}

Moreover, these findings are consistent with a previous study of Al Abdulwahab et al.⁴⁰ that concluded that

excessive body weight has a significant negative effect on the control of the muscles around the abdomen and back in the adult population.

These results are also consistent with Mayer et al.,⁴¹ who state that obesity, as measured by BMI, is correlated with decreased trunk muscular endurance. Duncan et al.⁴² also support these findings by reporting that overweight and obesity in British children is significantly associated with poor body function compared to children with normal weight.

Several explanations may clarify the decreased trunk muscular strength in overweight and obese children and adolescents. First, the possible variations between normal-weight adolescents and overweight and obese adolescents in the morphology and geometry of the trunk muscles could be due to excessive accumulation of adipose tissue inside and over the back muscles. This is in agreement with the hypothesis of Ryan et al.⁴³ who state that overweight and obese children may accumulate adipose tissue in the abdominal region, where the greatest fatty deposition occurs in the rectus abdominal muscle, followed by the lateral abdominal muscle and the erector spinae. This accumulation of adipose tissue inside the muscle may interfere with its functional capability, as Hicks et al.⁴⁴ have reported. The reduction of the functional capability of the core musculature, especially in terms of strength and endurance, may alter the induction of these muscles into the spinal stability.⁴⁵

Second, the possible variations between normal-weight adolescents and overweight and obese ones are in their general fitness levels. Most overweight and obese adolescents have lower fitness levels than normal-weight adolescents; although their extremities are strong, their trunks are weak. The trunk muscles fail to stabilise the body to enable the extremities to generate sufficient force to produce efficient movement patterns. This is in line with the results of Tse et al.⁴⁶ who find that core stability is essential to enable the trunk to control the movement of the distal extremities during participation in sports.

Third, weight is also an important factor of performance on measuring strength.^{47,48} On physical assessments demanding thrusting or elevating body mass, young obese persons show poor performance compared to non-obese young persons because of the effect of overload imposed by excess fat in addition to the original physical load.

Fourth, according to Cavuoto⁴⁹ and Maffiuletti et al.,⁵⁰ muscle fatigue in overweight and obese children negatively affects the strength and endurance periods during exercises. Moreover, overweight and obese children are not familiar with physical fitness and experience greater harassment during physical activities than normal-weight children.⁵¹

The strengths of the current study are that it included a large sample of adolescents and that it assessed the relationship between trunk flexors and extensors and BMI in paediatrics, which has not been extensively covered in previous studies.

The current study has some limitations that should be taken into consideration. First, the BMI measured in this study is the most popular index of adiposity, but it does not differentiate between fat and lean body masses. Second, only males participated in this study as access to female schools was not available.

Conclusion

This study clearly indicates that obesity is correlated with decreased trunk muscle torque and power in adolescents. Therapeutic interventions for overweight and obese adolescents should be supported by exercise training programmes to improve muscle performance variables including peak torque and power. These interventions and programmes would consequently enhance the quality of life of overweight and obese adolescents.

Recommendations

Further studies are required that examine the differences between males and females in trunk muscle performance across different age groups. Future studies should also examine the predictors of the trunk flexor and extensor torque using sociodemographic and clinical (BMI) variables.

Source of funding

This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Conflict of interest

The authors do not have any conflict of interest to declare.

Ethical approval

The Ethics Committee of College of Medical Rehabilitation Sciences, Taibah University, approved this study on 05/03/2020 (Approval No. CMR-PT-2020-03).

Consent

During the research activities, each student was informed about the study objectives, emphasising the confidentiality of collected data, and verbal consent was provided by the subject prior to the study.

Authors' contributions

Design and conceptualisation of the study, in addition to data analysis, were carried out by AMA and HAE. Data interpretation was done by all the authors. AMA and HAE conducted data collection and organisation procedures and provided the manuscript in its initial form. Critical review, modifications, and approval of the manuscript's final form were done by OAK and FSA. All authors conducted a similarity check using iThenticate software to approve the final form of the manuscript.

Acknowledgment

The authors would like to thank all participants for their efforts and patience during data collection.

References

- Cecily L, Betz. Special issue on childhood overweight and obesity. *J Pediatr Nurs* 2014; 29: 491–492.
- Al-Hussaini A, Bashir MS, Khormi M, AlTuraiki M, Alkhamis W, Alrajhi M, et al. Overweight and obesity among Saudi children and adolescents: where do we stand today? *Saudi J Gastroenterol* 2019; 25(4): 229. official journal of the Saudi Gastroenterology Association.
- Reilly JJ, Kelly J. Long-term impact of overweight and obesity in childhood and adolescence on morbidity and premature mortality in adulthood: systematic review. *Int J Obes* 2011; 35: 891–898.
- Wilson Louise F, Green Adele C, Jordan Susan J, Neale Rachel E, Webb Penelope M, Whiteman David C. The proportion of cancers attributable to social deprivation: a population-based analysis of Australian health data. *Cancer Epidemiology* 2020; 67: 101742.
- Silva MR, Badaró AF, Dall'Agnol MM. Low back pain in adolescent and associated factors: a cross sectional study with schoolchildren. *Braz J Phys Ther* 2014; 18(5): 402–409.
- Lafortuna CL, Maffiuletti NA, Agosti F, Sartorio A. Gender variations of body composition, muscle strength and power output in morbid obesity. *Int J Obes* 2005; 29(7): 833–841.
- Tomlinson DJ, Erskine RM, Morse CI, Winwood K, Onambélé-Pearson GL. Combined effects of body composition and ageing on joint torque, muscle activation and co-contraction in sedentary women. *Age* 2014; 36(3): 9652.
- Maffiuletti NA, Ratel S, Sartorio A, Martin V. The impact of obesity on in vivo human skeletal muscle function. *Curr Obes Rep* 2013; 2: 251–260.
- Delmonico MJ, Harris TB, Visser M, Park SW, Conroy MB, Velasquez-Mieyer P, et al. Longitudinal study of muscle strength, quality, and adipose tissue infiltration. *Am J Clin Nutr* 2009; 90(6): 1579–1585.
- Hilton Tiffany N, et al. Excessive adipose tissue infiltration in skeletal muscle in individuals with obesity, diabetes mellitus, and peripheral neuropathy: association with performance and function. *Phys Ther* 2008; 88(11): 1336–1344.
- Hulens M, Vansant G, Lysens R, Claessens AL, Muls E, Brumagne S. Study of differences in peripheral muscle strength of lean versus obese women: an allometric approach. *Int J Obes Relat Metab Disord* 2001; 25: 676–681.
- Pajoutan Mojdeh, Mehta Ranjana K, Cavuoto Lora A. Obesity effect on isometric strength of the trunk extensors. In: *Proceedings of the human factors and ergonomics society annual meeting*. Los Angeles, CA: SAGE Publications; 2016. 60 (1). Sage CA.
- Harreby M, Neergaard K, Hesselsoe G, Kjer J. Are radiologic changes in the thoracic and lumbar spine of adolescents risk factors for low back pain in adults? A 25-year prospective cohort study of 640 schoolchildren. *Spine* 1995; 20: 2298–2302.
- Leboeuf-Yde C, Kyvik KO. At what age does low back pain become a common problem? A study of 29,424 individuals aged 12–41 years. *Spine* 1998; 23: 228–234.
- Grabiner MD, Jeziorowski JJ, Divekar AD. Isokinetic measurements of trunk extension and flexion performance collected with the biodex clinical data station. *J Orthop Sports Phys Ther* 1990; 11(12): 590–598.
- Newton M, Thow M, Somerville D, Henderson I, Waddell G. Trunk strength testing with iso-machines. Part 2: experimental evaluation of the Cybex II Back Testing System in normal subjects and patients with chronic low back pain. *Spine* 1993; 18(7): 812–824.
- Iwai K, Nakazato K, Irie K, Fujimoto H, Nakajima H. Trunk muscle strength and disability level of low back pain in collegiate wrestlers. *Med Sci Sports Exerc* 2004; 36(8): 1296–1300.
- Yahia A, Jribi S, Ghroubi S, Elleuch M, Baklouti S, Habib Elleuch M. Evaluation of the posture and muscular strength of the trunk and inferior members of patients with chronic lumbar pain. *Joint Bone Spine* 2011; 78(3): 291–297.
- Den Hartog D, Eker HH, Tuinebreijer WE, Kleinrensink GJ, Stam HJ, Lange JF. Isokinetic strength of the trunk flexor muscles after surgical repair for incisional hernia. *Hernia* 2010; 14(3): 243–247.
- Hupli M, Sainio P, Hurri H, Alaranta H. Comparison of trunk strength measurements between two different isokinetic devices used at clinical settings. *J Spinal Disord* 1997; 10(5): 391–397.
- Langrana NA, Lee CK, Alexander H, Mayott CW. Quantitative assessment of back strength using isokinetic testing. *Spine* 1984; 9(3): 287–290.
- Brady S, Mayer T, Gatchel RJ. Physical progress and residual impairment quantification after functional restoration. Part II: isokinetic trunk strength. *Spine* 1994; 19(4): 395–400.
- Granacher U, Gollhofer A, Hortobagyi T, Kressig RW, Muehlbauer T. The importance of trunk muscle strength for balance, functional performance, and fall prevention in seniors: a systematic review. *Sports Med* 2013; 43(7): 627–641.
- Estrázulas JA, de Jesus K, da Silva RA, Libardoni Dos Santos JO. Evaluation isometric and isokinetic of trunk flexor and extensor muscles with isokinetic dynamometer: a systematic review. *Phys Ther Sport* 2020; 45: 93–102.
- Grabiner MD, Jeziorowski JJ. Isokinetic trunk extension discriminates uninjured subjects from subjects with previous low back pain. *Clin BioMech* 1992; 7(4): 195–200.
- García-Vaquero MP, Barbado D, Juan-Recio C, Lopez-Valenciano A, Vera- García FJ. Isokinetic trunk flexion extension protocol to assess trunk muscle strength and endurance: reliability, learning effect, and sex differences. *J Sport Health Sci* 2016; 328: 1–10.
- Juan-Recio C, Lopez-Plaza D, Barbado Murillo D, García-Vaquero MP, Vera- García FJ. Reliability assessment and correlation analysis of 3 protocols to measure trunk muscle strength and endurance. *J Sports Sci* 2018; 36(4): 357–364.
- Verbrugge J, Agten A, Eijnde BO, Vandenabeele F, De Baets L, Huybrechts X, et al. Reliability and agreement of isometric functional trunk and isolated lumbar strength assessment in healthy persons and persons with chronic nonspecific low back pain. *Phys Ther Sport* 2019; 38: 1–7.
- Roth R, Donath L, Kurz E, Zahner L, Faude O. Absolute and relative reliability of isokinetic and isometric trunk strength testing using the IsoMed- 2000 dynamometer. *Phys Ther Sport* 2017; 24: 26–31.
- Harding AT, Weeks BK, Horan SA, Little A, Watson SL, Beck BR. Validity and test-retest reliability of a novel simple back extensor muscle strength test (2017) *SAGE open medicine* 2017; 5: 1–9.
- Guilhem G, Giroux C, Couturier A, Maffiuletti NA. Validity of trunk extensor and flexor torque measurements using isokinetic dynamometry (2014) *J Electromyogr Kinesiol* 2014; 24(6): 986–993.
- Balague F, Nordin M, Skovron ML, Dutoit G, Yee A, Waldburger M. Non-specific low-back pain among schoolchildren: a field survey with analysis of some associated factors. *J Spinal Disord* 1994; 7(5): 374–379.
- Ogden CL, Flegal KM. Changes in terminology for childhood overweight and obesity. *Natl Health Stat Rep* 2010; 25: 1–5.
- Volek JS, Kraemer WJ, Bush JA, Boetes M, Incledon T, Clark KL, et al. Creatine supplementation enhances muscular

- performance during high-intensity resistance exercise. **J Am Dietetic Assoc** 1997; 97(7): 765–770.
35. Grabiner M, Jeziorowski J. Isokinetic trunk extension and flexion strength-endurance relationships. **Clin Biomech** 1991; 6: 118–122.
 36. Morini S, Ciccarelli A, Cerulli C, Giombini A, Di Cesare A, Ripani M. Functional anatomy of trunk flexion-extension in isokinetic exercise: muscle activity in standing and seated positions. **J Sports Med Phys Fit** 2008; 48: 17–23.
 37. Alcazar Julian, Csapo Robert, Ara Ignacio, Alegre Luis M. On the shape of the force-velocity relationship in skeletal muscles: the linear, the hyperbolic, and the double-hyperbolic. **Front Physiol** 2019; 10: 769.
 38. Maffiuletti NA, Jubeau M, Munzinger U, Bizzini M, Agosti F, De Col A, et al. Differences in quadriceps muscle strength and fatigue between lean and obese subjects. **Eur J Appl Physiol** 2007; 101: 51–59.
 39. Abdelmoula A, Martin V, Bouchant A, Walrand S, Lavet C, Taillardat M, et al. Knee extension strength in obese and nonobese male adolescents. **Appl Physiol Nutr Metabol** 2012; 37: 269–275.
 40. Alabdulwahab Sami S, Kachanathu Shaji John. Effects of body mass index on foot posture alignment and core stability in a healthy adult population. **J Exercise Rehabil** 2016; 12(3): 182.
 41. Mayer John M, Nuzzo James L, Chen Ren, Quillen William S, Verna Joe L, Miro Rebecca, et al. The impact of obesity on back and core muscular endurance in firefighters. **J Obesity** 2012; 1–7.
 42. Duncan MJ, Stanley M, Wright SL. The association between functional movement and overweight and obesity in British primary school children. **Sports Med Arthrosc Rehabil Ther Technol** 2013; 5(1): 11.
 43. Ryan AS, Harduarsingh-Permaul AS. Effects of weight loss and exercise on trunk muscle composition in older women. **Clin Interv Aging** 2014; 9: 395–402.
 44. Dederig A, Németh G, Harms-Ringdahl K. Correlation between electromyographic spectral changes and subjective assessment of lumbar muscle fatigue in subjects without pain from the lower back. **Clin BioMech** 1999; 14(2): 103–111.
 45. Hibbs AE, Thompson KG, French D, Wrigley A, Spears I. Optimizing performance by improving core stability and core strength. **Sports Med** 2008; 38(12): 995–1008.
 46. Tse MA, McManus AM, Masters RSW. Development and validation of a core endurance intervention program: implications for performance in college-age rowers. **J Strength Condit Res** 2005; 19(3): 547–552.
 47. Dumith SC, Ramires VV, Souza MA, Moraes DS, Petry FG, Oliveira ES, et al. Overweight/obesity and physical fitness among children and adolescents. **J Phys Activ Health** 2010; 7(5): 641–648 [PubMed: 20864760].
 48. Deforche B, Lefevre J, De Bourdeaudhuij I, Hills AP, Duquet W, Bouckaert J. Physical fitness and physical activity in obese and nonobese Flemish youth. **Obes Res** 2003; 11(3): 434–441 [PubMed: 12634442].
 49. Cavuoto Lora A, Maury A, Nussbaum. Differences in functional performance of the shoulder musculature with obesity and aging. **Int J Ind Ergon** 2013; 43(5): 393–399.
 50. Maffiuletti NA, Jubeau M, Munzinger U, Bizzini M, Agosti F, De Col A, et al. Differences in quadriceps muscle strength and fatigue between lean and obese subjects. **Eur J Appl Physiol** 2007; 101(1): 51–59.
 51. Sun Q, Van Dam RM, Spiegelman D, Heymsfield SB, Willett WC, Hu FB. Comparison of dual-energy X-ray absorptiometric and anthropometric measures of adiposity in relation to adiposity-related biologic factors. **Am J Epidemiol** 2010; 172(2): 1442–1454.

How to cite this article: Al-Shenqiti AM, Emara HA, Algarni FS, Khaled OA. Isokinetic trunk muscle performance in adolescents with different body mass indices. *J Taibah Univ Med Sc* 2021;16(4):550–557.