



Original Article

## Extensive drug-resistance in strains of *Escherichia coli* and *Klebsiella pneumoniae* isolated from paediatric urinary tract infections

Zakia Iqbal, M.Phil., Muhammad Z. Mumtaz, Ph.D.\* and Arif Malik, Ph.D.

Institute of Molecular Biology and Biotechnology, The University of Lahore, Lahore, Pakistan

Received 9 November 2020; revised 6 March 2021; accepted 10 March 2021; Available online 12 April 2021



### المخلص

**أهداف البحث:** تتزايد التهابات المسالك البولية في الأطفال بسرعة في جميع أنحاء العالم، وعادة ما تكون ناجمة عن البكتيريا المقاومة للأدوية. هذه الدراسة تحدد انتشار التهاب المسالك البولية في الأطفال، وتقيم نمط البكتيريا المقاومة للأدوية من قبل سلالات معزولة من الإشريكية والكليسيلا عن التهابات المسالك البولية في مرضى الأطفال.

**طرق البحث:** تم عزل السلالات البكتيرية الممرضة للجهاز البولي من المرضى الأطفال بالتهابات المسالك البولية التي تم إدخالها في معهد صحة الطفل بلاهور، باكستان. تم تحديد السلالات من جنس الإشريكية والكليسيلا، من خلال التوصيف الكيميائي الحيوي، وتعرضت لمقاييس الحساسية للمضادات الحيوية لتحديد البكتيريا المقاومة للأدوية ضد 21 من الأدوية المضادة للميكروبات.

**النتائج:** خلال ستة أشهر، تم عزل ما مجموعه 63 من الإشريكية و37 من سلالات الكليسيلا من 130 طفلاً مصابين بالتهابات المسالك البولية. وأظهرت مقاييس حساسية المضادات الحيوية أن الإشريكية والكليسيلا لها مقاومة عالية ضد كراموكسيكلاف، وسيفروكسيم، وسيفيكسيم، وسيفوتاكسيم، وسيفتازيديم، وسيفترياكسون، وسبيروفلوكساسين، وحمض ناليديكسك، ونورفلوكساسين، وحمض بيبيديمك، وكوتريموكسيزول. ومع ذلك، فقد وجد أن للعقاقير المضادة للميكروبات بما في ذلك بوليميكسين-ب، وكبريتات الكولستين، والكلورامفينيكول، ونيتروفورانتون، وفوسفومايسين فاعلية كبرى للتحكم بالسلالات الممرضة للجهاز البولي من القولونية الإشريكية والكليسيلا الرئوية. وتم تحديد السلالات الخمس التي تعاني من أعلى مقاومة ضد المضادات الحيوية المختبرة على أنها سلالات من الإشريكية القولونية والكليسيلا الرئوية.

**الاستنتاجات:** أظهرت دراستنا أن الإشريكية القولونية والكليسيلا الرئوية كانت البكتيريا المهيمنة لمقاومة للأدوية من التهابات المسالك البولية المكتسبة من

المجتمع في مجموعتنا. وتم الوصول إلى أن هذه البكتيريا الممرضة للجهاز البولي غير مستجيبة للعقارات المستخدمة بشكل روتيني من المضادات الحيوية مثل بيتا اللاكتام، والبيريدوبيريدين، والكينولون، والفلوروكينولون. قد تكون هذه النتائج مفيدة للأطباء لتعزيز العلاج التجريبي لالتهابات المسالك البولية في الأطفال.

**الكلمات المفتاحية:** متعددة مقاومة للأدوية؛ طب الأطفال؛ عدوى المسالك البولية؛ تسلسل الجينات؛ الكليسيلا الرئوية

### Abstract

**Objectives:** Urinary tract infections (UTIs) in children are rapidly increasing worldwide and are commonly caused by extensively drug-resistant bacteria. This study determines the prevalence of UTIs in paediatric patients and evaluates the pattern of extensively drug-resistance in *Escherichia coli* and *Klebsiella pneumoniae* strains isolated from paediatric UTI patients.

**Methods:** Uropathogenic bacterial strains were isolated from paediatric patients with UTIs admitted to the Institute of Child Health, Lahore, Pakistan. Strains of both *E. coli* and *K. pneumoniae* were identified using biochemical characterisation and subjected to antibiotic susceptibility assays for 21 common antimicrobial drugs in order to determine their extensively drug-resistant profile.

**Results:** We isolated 63 *E. coli* and 37 *K. pneumoniae* strains from 130 paediatric patients with UTIs over a period of six months. The antibiotic susceptibility assays showed that both the *E. coli* and *K. pneumoniae* strains exhibited a high degree of resistance against co-amoxiclav, cefuroxime, cefixime, cefotaxime, ceftazidime, ceftriaxone, ciprofloxacin, nalidixic acid, norfloxacin, pefedemox acid, and co-trimoxazole. However, several of the antimicrobial agents, including polymyxin

\* Corresponding address: Institute of Molecular Biology and Biotechnology, The University of Lahore, Main Campus, Lahore 54000, Pakistan.

E-mail: zahidses@gmail.com (M.Z. Mumtaz)

Peer review under responsibility of Taibah University.



B, colistin sulphate, chloramphenicol, nitrofurantoin, and fosfomycin, were found to retain their antimicrobial activities against both pathogens. The five highest antibiotic resistant strains were identified as *E. coli* strains ZK9, ZK40, and ZK60 and *K. pneumoniae* ZK32 and ZK89 using 16S rRNA gene sequencing.

**Conclusion:** Our study demonstrates that *E. coli* and *K. pneumoniae* are the dominant extensively drug-resistant uropathogenic bacteria in community-acquired UTIs in our cohort. These uropathogens were found to be resistant to the majority of the routinely-used classes of  $\beta$ -lactams, pyridopyrimidines, quinolones, and fluoroquinolone antibiotics, and these findings may be useful for clinicians in their treatment of paediatric UTIs.

**Keywords:** 16S rRNA gene sequencing; *Klebsiella pneumoniae*; Multiple drug-resistance; Paediatric infections; Urinary tract infections

© 2021 Taibah University.

Production and hosting by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Urinary tract infections (UTIs) are among the most prevalent and serious infections in children.<sup>1</sup> They frequently affect the lower urinary tract and are commonly referred to as bladder infections. UTIs may also affect the kidneys and are commonly referred to as pyelonephritis UTIs. Failure to effectively treat these infections may result in renal scarring, hypertension, and end-stage renal failure.<sup>2,3</sup> Owing to the high number of asymptomatic UTIs, its diagnosis and treatment remain challenging in this population. The risk of UTIs increases in hospitalised children with complicated severe acute malnutrition,<sup>4</sup> and UTIs are often diagnosed in children who exhibit poor toilet and hygiene habits, which may result in increased incidence of genetic transfer between pathogens and/or blockage in the normal urine flow between the ureters and kidney. UTIs are most common in girls as their urethra is closer to the anus.<sup>5</sup> The incidence of UTIs in boys increases within the first three months of starting school, but then decreases over time.<sup>6</sup> These infections are also commonly associated with the incorrect use of prescribed medications. This means that the rapid diagnosis and treatment of UTIs in small children is critical in preventing the long-term morbidities associated with renal scarring, such as hypertension, toxemia in pregnancy, chronic kidney disease, and ultimately the need for renal transplantation.<sup>2,3</sup>

UTIs are caused by Gram-negative and Gram-positive bacterial pathogens commonly found in the gastrointestinal tract. Gram-negative bacteria, including *Citrobacter* spp., *Enterococcus* spp., *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, and *Serratia* spp., and Gram-positive bacteria, including *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, and *Streptococcus* spp., are the most commonly identified uropathogenic

bacteria in these UTIs.<sup>7,8</sup> Such uropathogenic bacteria demonstrate extensive resistance to third-generation cephalosporins, classifying most of these infections as multiple drug-resistant (MDR) or extensive drug resistant (XDR).<sup>9–12</sup> MDR infections demonstrate resistance to more than one antimicrobial agent, while XDR have resistance to almost all antimicrobial agents.<sup>12</sup> Recent developments in paediatric urology have highlighted growing concerns around the emergence of extensive antibiotic resistance in urinary pathogens as a result of improper use and overuse of antibiotics.<sup>13</sup> Patients with sepsis in the intensive care unit also demonstrate high levels of MDR bacteria and produce morbid urine.<sup>14,15</sup> The antimicrobial susceptibility patterns in most pathogens demonstrate some geographical variation, and these changes are critical to establishing effective antimicrobial policies in local healthcare units. The use of antibiotics has resulted in a 95% reduction in infectious diseases with high mortality rates.<sup>16</sup> Given their effectiveness in mitigating health risks, antibiotic consumption has continued to increase in modern society.

UTI-causing uropathogenic bacteria experience extensive antimicrobial resistance with most strains exhibiting some resistance to nalidixic acid (81%), trimethoprim/sulfamethoxazole (83%), amoxicillin (67%), co-trimoxazole (61%), cephalexin (43%), gentamicin (49%), and ciprofloxacin (46%).<sup>17</sup> The discovery of novel antibiotics could help promote efforts to combat MDR and XDR in bacteria; however, where resistance is mediated by changes in cell wall permeability, novel antibiotics may be insufficient to combat widespread resistance.<sup>18</sup> Reduced permeability causes a reduction in the intracellular concentration of antibiotics and may be the result of efflux pumps, which actively transport the antibiotics out of the cell.<sup>19,20</sup> Bacteria also produce a variety of antibiotic metabolising enzymes, including  $\beta$ -lactamases, aminoglycoside-modifying enzymes, and chloramphenicol acetyltransferases that inactivate the antibiotics before they can exert their effects. Variations in the target site might also occur, in which case the antibiotic will lose its efficacy.<sup>21</sup> Most of these genes are encoded on plasmids, which may be transferred between bacterial populations, resulting in the emergence of the ever-growing numbers of MDR strains. These gene transfers have been studied extensively, and remain a critical concern in antimicrobial resistance.<sup>22</sup> This study determines the prevalence of UTIs and the frequency of the most commonly-found MDR and XDR uropathogenic *E. coli* and *Klebsiella* spp. in paediatric patients from Lahore, Pakistan. These uropathogenic bacteria are characterised using biochemical assays and screened for antibiotic resistance before final identification using 16S rRNA gene sequencing.

## Materials and Methods

### Urine collection

This study was conducted using a cohort of paediatric patients being treated in the outpatient department (OPD) and inpatient departments (nephrology, neonatal unit, general medical, urology, and neurology wards) of The Children's Hospital and the Institute of Child Health in Lahore, Pakistan, between June 2018 and February 2019. Freshly

voided midstream urine specimens were collected from both male and female patients aged 0–13 years suffering from high fever, abdominal pain, and vomiting. Urine samples were collected using a clean catch method in dry, sterile, wide neck, leak-proof containers. Informed consent from each participant or their guardians was recorded on a form that included explicit consent for the collection, storage, and testing of the samples. More than 130 specimens were aseptically collected and transported to the Laboratory of Microbiology at the Institute for Molecular Biology and Biotechnology at the University of Lahore in Pakistan. The children were diagnosed with symptomatic UTI and presented with a variety of symptoms, including fever, chills, nausea, vomiting, dysuria, frequency, urgency, incontinence, and flank pain.<sup>23</sup> The 5 mL urine samples were centrifuged at 10,000×g for 5 min and the supernatant was stained and processed to estimate white blood cell counts (pus cells) using a light microscope at low (10x) and high (40x) power as described by Cappuccino and Sherman.<sup>24</sup> Urine samples with a white blood cell count [WBC] of ≥5 cells/high power field (HPF) were considered positive for UTI and were then used to isolate bacterial pathogens.

#### Isolation of uropathogenic bacteria

Uropathogenic bacteria were isolated from urine samples using cysteine lactose electrolyte-deficient (CLED) agar.<sup>25</sup> Urine was poured over autoclaved CLED agar media at different dilutions and incubated at 37 °C for 72 h. Colonies with a yellow, opaque, or slightly deeper colour were tentatively identified as *Escherichia* spp., while large mucoid yellow or yellow-white colonies were considered to belong to the *Klebsiella* spp. Selected colonies were then picked, purified, and preserved at –20 °C until further experiments.

#### Biochemical characterization

The isolated strains were screened for Gram staining using the Duguid et al. method.<sup>26</sup> To differentiate between lactose fermenting and non-fermenting bacterial strains, freshly grown strains were streaked on MacConkey agar and incubated at 37 °C for 24 h. After incubation, bacterial strains were evaluated for pigmentation and any colony appearing as pink-red was classified as a lactose fermenting organism. To check the haemolysis of each bacterial strain, freshly grown strains were streaked on blood agar plates and incubated at 37 °C for 24 h. A clear zone around the colony confirmed the haemolytic capacity of each of the screened strains and enzymatic activities, such as oxidase, catalase, and urease activities, were then evaluated. Strains were also evaluated for citrate utilisation, motility, and indole production using biochemical tests, as described by Cappuccino and Sherman.<sup>24</sup>

#### Antibiotic resistance test

The disk diffusion method, M100, from the Clinical and Laboratory Standard Institute (CLSI)<sup>27</sup> was used to evaluate the antimicrobial susceptibility of each strain against 21 different antibiotics, including co-amoxiclav, amikacin, cefuroxime, cefixime, cefotaxime, ceftazidime, ceftriaxone,

ciprofloxacin, nalidixic acid, nitrofurantoin, norfloxacin, piperimic acid, sulbactam/cefoperazone, colistin sulphate, fosfomicin, meropenem, imipenem, chloramphenicol, piperacillin/tazobactam, co-trimoxazole, and polymyxin B. Uropathogenic strains were freshly grown overnight and heavily streaked on Mueller-Hinton (MH) plates.<sup>28</sup> Then, commercially available 4 mm antibiotic paper discs (Abtek Biologicals Ltd, UK) were placed on the MH plates and incubated at 37 °C for 24 h. Subsequently, the zone of inhibition in the bacterial lawn was then measured and recorded. *E. coli* ATCC 23509 and *E. coli* ATCC 25922 were also tested for antibiotic susceptibility and used as the control for our observations. The diameters of the zones of inhibition were then categorised as sensitive or resistant using the standard diameters provided by the CLSI.<sup>27</sup> The non-susceptible strains were classified as MDR, XDR, and pan drug-resistance (PDR) based on the criteria reported in Mogiorakos et al.<sup>12</sup> Non-susceptible strains presenting with resistance to ≥1 agent were non-susceptible; strains resistant to ≥3 antimicrobial categories were considered MDR, while those with susceptibility to ≥1 agent in all but ≤2 antimicrobial categories were considered to be XDR. Those bacterial strains that were resistant to all the listed antimicrobial agents were considered PDR.

#### Identification of uropathogenic bacterial strains using 16S rRNA gene sequencing

The genomic DNA from selected antibiotic-resistant uropathogenic bacterial strains (ZK9, ZK32, ZK40, ZK60, and ZK89) was isolated using proteinase K treatment and the polymerase chain reaction was performed as described by Mumtaz et al.<sup>29</sup> The PCR product was then sequenced using a commercial service provided by Macrogen (Seoul, Korea) and the sequences were identified using the BLAST algorithm from the NCBI web browser. The phylogenetic tree of closely related species was constructed using MEGA 7.0.26. DNA sequences were aligned using ClustalW, and phylogeny was determined using the maximum parsimony method.

## Results

Uropathogenic *Escherichia* and *Klebsiella* spp. were isolated from urine samples collected from 130 paediatric UTI patients in various wards, including OPD (29), nephrology (23), neonatal unit (18), general medical (12), urology (11), and neurology (7), of the Children's Hospital and Institute of Child Health in Lahore, Pakistan (Table 1). The

**Table 1: Distribution of selected paediatric patients across different hospital wards.**

Wards	n (%)
Out Patient Department	29
Nephrology	23
Neonatal unit	18
General medical	12
Urology	11
Neurology	7

**Table 2: Baseline demographic and clinical characterization of paediatric UTI patients.**

Variable	n (%)
<b>Age</b>	
Less than 1 year	23
1 year to 14 years	77
<b>Gender</b>	
Male	51
Female	49
<b>Vomiting</b>	
Yes	64
No	36
<b>Fever</b>	
Yes	87
No	13
<b>Complications</b>	
Immunosuppression	11
Urethra malformation	34
Neurological disorders	7
<b>UTI history</b>	
Yes	26
No	74
<b>Bacterial count</b>	
> 10 <sup>3</sup> CFU mL <sup>-1</sup>	28
> 10 <sup>5</sup> CFU mL <sup>-1</sup>	72

demographic data of these paediatric patients are summarised in Table 2. Some urine samples (30) exhibited no bacterial growth on the CLED agar and were thus excluded from the study. Of the remaining 100 paediatric patients, 23 were less than one year old and 77 were in the range of 1–14 years of age. UTI prevalence between the two sexes revealed that 51% of the UTIs were found in male children and 49% in female paediatric patients. Uropathogens were isolated from both symptomatic UTIs, presenting with vomiting (64%), fever (87%), and non-symptomatic UTIs, without vomiting (36%) or fever (13%). Health complications, including immunosuppression (11%), urethral malformation (34%), and neurological disorders (7%), were also identified in several UTI patients. A total of 26% of the paediatric patients had a history of


antibiotic use and the highest bacterial count, 10<sup>5</sup> CFU mL<sup>-1</sup>, was observed in 72% of urine samples, while 28% of the samples had a reduced bacterial cell count of 10<sup>3</sup> CFU mL<sup>-1</sup> (Table 2).

#### Characterization of the uropathogenic strains

A total of 100 strains were isolated and identified as *Escherichia* and *Klebsiella* spp. using morphological and biochemical characterisation. These isolates were coded as ZK1, ZK2, ZK3, ... to ZK100 and divided into groups A and B based on the results of the biochemical characterisation (Table 3). Biochemical characterisation of selected strains revealed that all the strains in both groups were Gram-negative, positive for catalase activity, and negative for oxidase activities. All the strains presented with a yellow acidic slant and butt with gas production in the TSI test. Group A strains were negative for urease activity and citrate utilisation tests but positive for indole production and motility. The group A strains were negative for urease activity and citrate utilisation, however, group B strains were positive to these assays (Table 3). Based on the biochemical characterisation, group A strains were identified as belonging to the *Escherichia* spp. and group B strains were identified as *Klebsiella* spp. (Table 3 and Figure 1a). Among the *E. coli* strains, 38% were isolated from male patients and 25% were from female patients. In the *Klebsiella* strains, 21% of the strains were from boy and 16% were from girl patients (Figure 1b).

#### Antibiotic susceptibility

The susceptibility patterns of the Gram-negative uropathogens is summarized in Figures 2 and 3. Overall, ≥90% of the uropathogenic strains were not susceptible to cefuroxime, cefixime, and cefotaxime. More than 80% of uropathogenic strains were also found to be resistant to co-amoxiclav, ceftazidime, ceftriaxone, ciprofloxacin, nalidixic acid, norfloxacin, piperidic acid, and co-trimoxazole. The antimicrobial susceptibility patterns of the *Escherichia* and *Klebsiella* spp. are shown in Figures 2 and 3. All of the *E. coli* strains (group A) were screened against all 21 antimicrobial



**Figure 1:** Prevalence of uropathogenic *Escherichia* and *Klebsiella* strains (A) and their distribution in male and female paediatric UTIs (B).




Figure 2: Antimicrobial susceptibility of uropathogenic *Escherichia* strains.

Table 3: Characterization of the uropathogenic bacterial strains identified in paediatric UTI patients.

Characterization	Group A (n = 63)*	Group B (n = 37)**
Gram-staining	Gram-negative	Gram-negative
Catalase activity	Positive	Positive
Oxidase activity	Negative	Negative
Urease activity	Negative	Positive
Citrate utilization test	Negative	Positive

\* Group A was identified as *Escherichia* spp. through biochemical characterization. \*\* Group B was identified as *Klebsiella* spp. through biochemical characterization.

drugs, and the results are depicted in Figure 2. The highest degree of resistance (93%) was reported for cefixime, followed by cefuroxime (90%). More than 80% of the *E. coli* strains were not susceptible to cefotaxime, ceftazidime, ceftriaxone, ciprofloxacin, nalidixic acid, norfloxacin, pefedemic acid, and co-trimoxazole. However, *E. coli* strains were susceptible to colistin sulphate (85%), polymyxin B (85%), chloramphenicol (84%), nitrofurantoin (79%), fosfomycin (76%), and meropenem (71%). The *Klebsiella* strains (group B) demonstrated a similar pattern of resistance and susceptibility (Figure 3). The highest





Figure 3: Antimicrobial susceptibility of uropathogenic *Klebsiella* strains.

**Table 4: Multiple drug-resistant and extensively drug-resistant uropathogenic *E. coli* and *K. pneumoniae* from paediatric UTIs in Lahore, Pakistan.**

Antimicrobial resistance	Uropathogen identifier
Multiple drug-resistant (MDR)	<i>Escherichia</i> spp. strains (n = 5) ZK52, ZK53, ZK63, ZK64, ZK98, <i>Klebsiella</i> spp. strains (n = 9) ZK19, ZK20, ZK42, ZK46, ZK47, ZK54, ZK55, ZK97, ZK99
Extensively drug resistant (XDR)	<i>Escherichia</i> spp. strains (n = 58) ZK1, ZK3, ZK4, ZK6, ZK7, ZK9 <sup>a</sup> , ZK10, ZK11, ZK12, ZK13, ZK16, ZK21, ZK22, ZK23, ZK24, ZK26, ZK27, ZK29, ZK31, ZK33, ZK34, ZK38, ZK39, ZK40 <sup>a</sup> , ZK44, ZK49, ZK56, ZK57, ZK58, ZK59, ZK60 <sup>a</sup> , ZK61, ZK65, ZK66, ZK67, ZK68, ZK69, ZK70, ZK71, ZK73, ZK76, ZK78, ZK79, ZK80, ZK81, ZK82, ZK83, ZK84, ZK85, ZK87, ZK88, ZK90, ZK91, ZK92, ZK93, ZK95, ZK96, ZK100 <i>Klebsiella</i> spp. strains (n = 28) ZK2, ZK5, ZK8, ZK14, ZK15, ZK17, ZK18, ZK25, ZK28, ZK30, ZK32 <sup>a</sup> , ZK35, ZK36, ZK37, ZK41, ZK43, ZK45, ZK48, ZK50, ZK51, ZK62, ZK72, ZK74, ZK75, ZK77, ZK86, ZK89 <sup>a</sup> , ZK94,

<sup>a</sup> Indicated XDR strains were selected for molecular identification using 16S rRNA gene sequencing.



**Figure 4:** Phylogenetic tree describing the 16S rRNA gene sequences of *E. coli* strains ZK9, ZK40, ZK60 (accession number MT764342, MT764344, and MT764345, respectively) and *K. pneumoniae* strains ZK32 and ZK89 (accession number MT764343 and MT764346, respectively) and the other closely related bacterial strains found in the GenBank database.

resistance (95%) was observed against cefixime, followed by cefuroxime (92%) and cefotaxime (92%). More than 75% of the *Klebsiella* strains were not susceptible to co-amoxiclav, ceftazidime, ceftriaxone, ciprofloxacin, nalidixic acid, norfloxacin, pefedemic acid, and co-trimoxazole. Antimicrobial agents polymyxin B (89%), colistin sulphate (86%), chloramphenicol (81%), nitrofurantoin (78%), and fosfomycin (76%) were more effective against the *Klebsiella* strains (Figure 3). The estimated MDR and XDR patterns are reported in Table 4 and reveal that 14% of the strains can be classified as MDR, and 86% of the uropathogenic strains identified in this study could be classified as XDR. None of the strains were found to be PDR. Among the MDR strains, five were identified as *Escherichia* spp. and nine were *Klebsiella* spp. In contrast, 58 of the *Escherichia* spp. and 28 of the *Klebsiella* spp. were found to be XDR. The XDR strains ZK9, ZK32, ZK40, ZK60, and ZK89 were shown to demonstrate the highest degree of resistance to the antibiotics used in this study and reported resistance to 17 of the 21 antimicrobials evaluated here. These strains were then selected for molecular identification.

#### Identification of selected antibiotic-resistant uropathogens

The five most representative MDR strains, ZK9, ZK32, ZK40, ZK60, and ZK89, were selected for molecular identification using 16S rRNA partial gene sequencing. These strains were identified as *Escherichia coli* strain ZK9, *Klebsiella pneumoniae* strain ZK32, *Escherichia coli* strain ZK40, *Escherichia coli* strain ZK60, and *Klebsiella pneumoniae* strain ZK89. A phylogenetic tree of these strains was constructed using the neighbour-joining method and broadened by selecting closely matched *Enterobacteriaceae* strains, as shown in Figure 4. The resulting sequences were deposited in NCBI GenBank with accession numbers MT764342 for ZK9, MT764343 for ZK32, MT764344 for ZK40, MT764345 for ZK60, and MT764346 for ZK89 (Figure 4).

#### Discussion

Antimicrobial drug resistance is a life-threatening issue, especially in paediatric patients, when treating infections caused by *Escherichia* and *Klebsiella* spp. Antimicrobial drug-resistance is increasing every day and the normal microbial flora in the urinary tract is becoming a reservoir for resistance genes.<sup>30</sup> In this study, hundreds of uropathogenic *Escherichia* and *Klebsiella* strains were isolated from paediatric patients with UTIs and their antimicrobial susceptibility was evaluated. The *E. coli* and *K. pneumoniae* strains isolated from the paediatric patients in this study showed the highest antimicrobial resistance ( $\geq 60\%$ ) against tested antibiotics. Among these strains,  $\geq 80\%$  were found to be not susceptible to cefixime, cefuroxime, cefotaxime, nalidixic acid, ceftazidime, co-trimoxazole, pefedemic acid, ceftriaxone, ciprofloxacin, and norfloxacin, and 86% of these strains, 58 *Escherichia* strains and 28 *Klebsiella* strains, were found to be XDR. Similar findings have also been reported in various clinical settings across Nepal and Ethiopia.<sup>9,31,32</sup> However, the prevalence of MDR *Escherichia* and *Klebsiella* strains has not yet been reported for Lahore, Pakistan. This study addresses this lack of

information and provides insight into the increasing incidence of MDR in paediatric UTIs in Pakistan.

In this study, uropathogenic bacterial strains were identified as *E. coli* and *K. pneumoniae* based on their biochemical characterisation. These uropathogens are Gram-negative and are known to be the dominant pathogens in UTIs across the world due to their unique structure, which facilitates their attachment to the host cells.<sup>33</sup> These structural features help the bacteria avoid exclusion by urinary lavage and support bacterial reproduction and expansion in these tissues, supporting invasive infection and pyelonephritis.<sup>34</sup> The *Escherichia* strains in this study were found to be negative for citrate utilisation and positive for indole production, while the *Klebsiella* strains showed the opposite result. Indole and citrate-utilization tests are commonly used to identify the *Enterobacteriaceae* including *Escherichia*, *Enterobacter*, and *Klebsiella* spp.<sup>35</sup> In this study the *Escherichia* strains were indole positive, which is likely related to their ability to produce tryptophanase enzymes that can convert tryptophan into indole. When indole reacts with para-dimethyl-amino benzaldehyde, a pink-coloured complex is produced that differentiates *Klebsiella* from *Enterobacter* spp. In this study, all *Escherichia* and *Klebsiella* strains were oxidase negative, which may be due to the inactivation of cytochrome C-oxidase and an inability to utilise oxygen for energy production via the electron transfer chain, which confirms their link to the *Enterobacteriaceae* family.<sup>36</sup> In our study, the production of a yellow slant and butt with gas and no H<sub>2</sub>S on TSI media indicates the carbohydrate fermentation ability of these strains.<sup>37</sup> However, the molecular identification method is more reliable and provides information up to the species level. In this study, the evaluation of the 16s rRNA gene revealed that most MDR strains were *Escherichia coli* strain ZK9, *Klebsiella pneumoniae* strain ZK32, *Escherichia coli* ZK40, *Escherichia coli* strain ZK60, and *Klebsiella pneumoniae* strain ZK89. Strains ZK9, ZK40, and ZK60 were shown to be closely related to *Escherichia* sp. strain UA32, which is an MDR strain isolated from female patients during pregnancy.<sup>38</sup> The XDR *Klebsiella pneumoniae* strains ZK32 and ZK89 were shown to be closely related to MDR *Klebsiella* sp. strain UA47 from pregnant females.<sup>38</sup>

In this study, the highest degree of resistance ( $\geq 90\%$ ) in the *E. coli* strains was observed for cefixime and cefuroxime, while the antimicrobial agents cefotaxime, ceftazidime, ceftriaxone, ciprofloxacin, nalidixic acid, norfloxacin, pefedemic acid, and co-trimoxazole reported a  $\geq 80\%$  non-susceptibility rate in these strains. Recently, Vazouras et al.<sup>39</sup> reported their findings from Greece, wherein 79% of the strains in their study were *E. coli*, making them the most common UTI-causing organism in paediatric patients. This study also reported the resistance rates for ampicillin (42%), trimethoprim/sulfamethoxazole (up to 26.5%), amoxicillin/clavulanic acid (up to 12.2%), cephalosporins (up to 1.7%), nitrofurantoin (up to 2.3%), ciprofloxacin (up to 1.4%), and amikacin (up to 0.9%), which were much lower than those found in our study. This difference may be the result of geographical variations resulting from local differences in the frequency of antibiotic use and ease of antibiotic availability.<sup>40</sup> Kayastha et al.<sup>9</sup> reported a rising prevalence in resistance, up to 62%, of extended-spectrum  $\beta$ -lactamase-

producing Gram-negative bacteria in paediatric UTIs and Shrestha et al.<sup>10</sup> reported MDR uropathogenic *E. coli* (53%), *E. faecalis* (22%), *K. pneumoniae* (7%), and *S. aureus* (7%). Hsueh et al.<sup>11</sup> evaluated urogenital pathogens and found that 23% of their urine samples were positive for the most common resistance factors in bacterial uropathogens. Mirzarazi et al.<sup>41</sup> reported that their *E. coli* isolates were resistant to nalidixic acid and trimethoprim-sulfamethoxazole. In this study, we found that the *Klebsiella* strains ( $\geq 92\%$ ) were not susceptible to cefixime, cefuroxime, or cefotaxime. More than 75% of these strains were also resistant to co-amoxiclav, ceftazidime, ceftriaxone, ciprofloxacin, nalidixic acid, norfloxacin, pefloxacin, and co-trimoxazole. Similarly, Hayat et al.<sup>42</sup> reported that the prevalence of paediatric UTIs was up to 52% in Pakistan, with the majority of infections occurring due to extended-spectrum  $\beta$ -lactamase-producing *K. pneumoniae*, which demonstrated a high degree of resistance (96%) to ceftazidime. *Klebsiella* spp. also showed resistance to trimethoprim-sulfamethoxazole, ciprofloxacin, and nalidixic acid.<sup>41</sup>

Across the world, most patients carry a variety of MDR genes that have developed as a result of the misuse and improper administration of antibiotics. This has resulted in the widespread sharing of  $\beta$ -lactamase enzymes in the commensal bacteria, promoting the development of novel MDR strains.<sup>43</sup> In our study, we observed the highest MDR patterns for the aminoglycosides and  $\beta$ -lactam antibiotics. This is of significant clinical concern, especially in children, as their immune system is still developing.<sup>44</sup> This is particularly problematic when uropathogenic organisms present with MDR characteristics transferred from other organisms via genetic exchange.<sup>45</sup> Longer hospital stays also result in infections with extended-spectrum  $\beta$ -lactamase-producing organisms.<sup>46</sup> MDR could also be due to an unavoidable genetic reaction to the solid discriminating strength forced by antibiotic therapy which plays its dynamic role in the development of MDR in uropathogens and their plasmid having MDR genes.<sup>25</sup>

## Conclusion

Our data shows that the majority of the uropathogenic *Escherichia* and *Klebsiella* strains isolated in this study are resistant to cefixime, cefuroxime, nalidixic acid, and cefotaxime, suggesting that such antibiotic treatment should be suspended following extensive long-term evaluations. Self-medication and improper diagnosis should be avoided to prevent increasing antibiotic resistance in both of these clinically-important bacterial genera.

## Recommendations

Uropathogenic *E. coli* and *K. pneumoniae* were found to be responsible for the majority of the MDR UTIs identified in paediatric patients in Lahore, Pakistan. Efforts must be made to control MDR infections in paediatric patients by executing evidence-based monitoring of UTI treatment and creating awareness of the judicious use of antibiotics.

## Source of funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## Conflict of interest

The authors have no conflict of interest to declare.

## Ethical approval

Ethical approval for this study protocol was granted by the Institute of Molecular Biology and Biotechnology at The University of Lahore, Pakistan, with the reference number IMBB-744 and dated 10-11-2018.

## Authors' contributions

ZI conducted the research, provided the research materials, and collected and organised the data. MZM conceptualised and supervised the study, analysed and interpreted the data, and wrote the initial and final drafts of the manuscript. AM provided the research materials and logistical support for this project. All authors critically reviewed and approved the final draft of this manuscript and are responsible for the content and similarity index of the manuscript.

## Acknowledgments

The authors would like to acknowledge the research facility provided by the Institute of Molecular Biology and Biotechnology (IMBB) at the University of Lahore (UOL), Lahore, Pakistan.

## References

1. Mahony M, McMullan B, Brown J, Kennedy SE. Multidrug-resistant organisms in urinary tract infections in children. *Pediatr Nephrol* 2020; 35(9): 1–11.
2. Lane DR, Takhar SS. Diagnosis and management of urinary tract infection and pyelonephritis. *Emerg Med Clin* 2011; 29(3): 539–552.
3. Whiting P, Westwood M, Watt I, Cooper J, Kleijnen J. Rapid tests and urine sampling techniques for the diagnosis of urinary tract infection (UTI) in children under five years: a systematic review. *BMC Pediatr* 2005; 5(1): 1–13.
4. Uwaezuoke SN, Ndu IK, Eze IC. The prevalence and risk of urinary tract infection in malnourished children: a systematic review and meta-analysis. *BMC Pediatr* 2019; 19(1): 1–20.
5. Stauffer CM, Weg B, Donadini R, Ramelli GP, Marchand S, Bianchetti MG. Family history and behavioral abnormalities in girls with recurrent urinary tract infections: a controlled study. *J Urol* 2004; 171(4): 1663–1665.
6. Rabasa AI, Shattima D. Urinary tract infection in severely malnourished children at the University of Maiduguri Teaching Hospital. *J Trop Pediatr* 2002; 48(6): 359–361.



7. Mickymaray S, Al Aboody MS. In vitro antioxidant and bactericidal efficacy of 15 common spices: novel therapeutics for urinary tract infections? *Medicina* **2019**; 55(6): 1–18.
8. Behzadi P, Urbán E, Matuz M, Benkő R, Gajdács M. The role of Gram-negative bacteria in urinary tract infections: current concepts and therapeutic options. In: *Advances in experimental medicine and biology*. New York: Springer; 2020.
9. Kayastha K, Dhungel B, Karki S, Adhikari B, Banjara MR, Rijal KR, et al. Extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* and *Klebsiella* species in pediatric patients visiting international friendship children's hospital, Kathmandu, Nepal. *Infect Dis Res Treat* **2020**; 13: 1–7. <https://doi.org/10.1177/1178633720909798>.
10. Shrestha LB, Baral R, Poudel P, Khanal B. Clinical, etiological and antimicrobial susceptibility profile of pediatric urinary tract infections in a tertiary care hospital of Nepal. *BMC Pediatr* **2019**; 19(1): 36.
11. Hsueh PR, Hoban DJ, Carmeli Y, Chen SY, Desikan S, Alejandria M, Binh TQ. Consensus review of the epidemiology and appropriate antimicrobial therapy of complicated urinary tract infections in Asia-Pacific region. *J Infect* **2011**; 63(2): 114–123.
12. Magiorakos AP, Srinivasan A, Carey RT, Carmeli Y, Falagas MT, Giske CT, et al. Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* **2012**; 18(3): 268–281.
13. Mojtahedzadeh M, Panahi Y, Fazeli MR, Najafi A, Pazouki M, Navehsi BM, Beiraghdar F. Intensive care unit-acquired urinary tract infections in patients admitted with sepsis: etiology, risk factors, and patterns of antimicrobial resistance. *Int J Infect Dis* **2008**; 12(3): 312–318.
14. World Health Organization (WHO). *Urinary tract infections in infants and children in developing countries in the context of IMCI (No. WHO/FCH/CAH/05.11)*; 2005.
15. Shepherd AK, Pottinger PS. Management of urinary tract infections in the era of increasing antimicrobial resistance. *Med Clin* **2013**; 97(4): 737–757.
16. Afsharpaiman S, Bairaghdar F, Torkaman M, Kavehmanesh Z, Amirsalari S, Moradi M, Safavimirmahalleh M. Bacterial pathogens and resistance patterns in children with community-acquired urinary tract infection: a cross sectional study. *J Compr Pediatr* **2012**; 3(1): 16–20.
17. Ray J, Paul R, Haldar A, Mondol S. A study on antibiotic resistance pattern of *Escherichia coli* isolated from urine specimens in Eastern India. *Int J Med Sci Publ Health* **2015**; 4(12): 1670–1674.
18. Ghai I, Ghai S. Understanding antibiotic resistance via outer membrane permeability. *Infect Drug Resist* **2018**; 11: 523–530.
19. Masi M, Réfregiers M, Pos KM, Pagès JM. Mechanisms of envelope permeability and antibiotic influx and efflux in Gram-negative bacteria. *Nature Microb* **2017**; 2(3): 1–7.
20. Džidic S, Šuškovac J, Kos B. Antibiotic resistance mechanisms in bacteria: biochemical and genetic aspects. *Food Technol Biotechnol* **2008**; 46: 11–21.
21. Dockrell HM, Goering RV, Roitt I, Wakelin D, Zuckerman M. In: Mims C, Dockrell HM, Goering RV, Roitt I, Wakelin D, Zuckerman M, editors. *Attacking the enemy: antimicrobial agents and chemotherapy. Medical Microbiology*. Netherlands: Elsevier; 2004. pp. 473–507.
22. Mills S, McAuliffe OE, Coffey A, Fitzgerald GF, Ross RP. Plasmids of lactococci-genetic accessories or genetic necessities? *FEMS Microbiol Rev* **2006**; 30: 243–273.
23. Woldemariam HK, Geleta DA, Tulu KD, Aber NA, Legese MH, Fenta GM, et al. Common uropathogens and their antibiotic susceptibility pattern among diabetic patients. *BMC Infect Dis* **2019**; 19: 1–10.
24. Cappuccino JC, Sherman N. *Microbiology*. 3<sup>rd</sup> ed. A Laboratory Manual; 1992. pp. 125–179.
25. Chakraborty SP, Mahapatra SK, Roy S. Biochemical characters and antibiotic susceptibility of *Staphylococcus aureus* isolates. *Asian Pac J Trop Biomed* **2011**; 1(3): 212–216.
26. Duguid JP. Staining methods. In: Collee JG, Fraser AG, Marmion BP, Simmons A, editors. *Mackie and McCartney practical medical microbiology*. Edinburgh: Churchill Livingstone; 1996.
27. CLSI. *Performance standards for antimicrobial susceptibility testing (M100)*. 30<sup>th</sup> ed. Weinstein, MP: Clinical and Laboratory Standards Institute (CLSI); 2020.
28. Murray PR, Zeiting JR. Evaluation of Mueller-Hinton agar for disk diffusion susceptibility tests. *J Clin Microbiol* **1983**; 18(5): 1269–1271.
29. Mumtaz MZ, Ahmad M, Jamil M, Hussain T. Zinc solubilizing *Bacillus* spp. potential candidates for biofortification in maize. *Microbiol Res* **2017**; 202: 51–60.
30. Okeke IN, Fayinka ST, Lamikanra A. Antibiotic resistance in *Escherichia coli* from Nigerian students. *Emerg Infect Dis* **2000**; 6(4): 393–396.
31. Shakya P, Shrestha D, Maharjan E, Sharma VK, Paudyal R. ESBL production among *E. coli* and *Klebsiella* spp. causing urinary tract infection: a hospital-based study. *Open Microbiol J* **2017**; 11: 23–30.
32. Seifu WD, Gebissa AD. Prevalence and antibiotic susceptibility of uropathogens from cases of urinary tract infections (UTI) in Shashemene referral hospital, Ethiopia. *BMC Infect Dis* **2018**; 18(1): 1–9.
33. Vaca DJ, Thibau A, Schütz M, Kraiczky P, Happonen L, Malmström J, et al. Interaction with the host: the role of fibronectin and extracellular matrix proteins in the adhesion of Gram-negative bacteria. *Med Microbiol Immunol* **2020**; 209(3): 277–299.
34. Ayelign B, Abebe B, Shibeshi A, Meshesha S, Shibabaw T, Addis Z, et al. Bacterial isolates and their antimicrobial susceptibility patterns among pediatric patients with urinary tract infections. *Turk J Urol* **2018**; 44(1): 62–69.
35. Bahjat SA, Al-Taeef MF, Al-Hassani AM, Al-Hassani OM. Molecular and bacteriological method for identification of lactose fermenting *Salmonella* in Mosul province. *Indian J Public Health Res Dev* **2019**; 10(12): 1428–1434.
36. Prescott LM, Harley JP, Klein DA. *Isolation of pure bacterial cultures from specimens: Microbiology*. Boston: McGraw-Hill Companies; 1999.
37. Buxton A, Fraser G. *Animal microbiology, vol. 1: immunology, bacteriology, mycology, diseases of fish and laboratory methods*. Blackwell Scientific Publications; 1977.
38. Asmat U, Mumtaz MZ, Malik A. Rising prevalence of multidrug-resistant uropathogenic bacteria from urinary tract infections in pregnant women. *J Taibah Univ Medical Sci* **2020**; 16(1): 102–111.
39. Vazouras K, Velali K, Tassiou I, Anastasiou-Katsiardani A, Athanasopoulou K, Barbouni A, et al. Antibiotic treatment and antimicrobial resistance in children with urinary tract infections. *J Glob Antimicrob Resist* **2020**; 20: 4–10.
40. Zorc JJ, Kiddoo DA, Shaw KN. Diagnosis and management of pediatric urinary tract infections. *Clin Microbiol Rev* **2005**; 18(2): 417–422.
41. Mirzarazi M, Rezaatofghi SE, Pourmahdi M, Mohajeri MR. Antibiotic resistance of isolated Gram-negative bacteria from urinary tract infections (UTIs) in Isfahan. *Jundishapur J Microbiol* **2013**; 6(8): 1–5.
42. Hayat S, Siddique MH, Aslam B, Nadeem H, Ashraf A, Saqalein M, et al. Extended-spectrum- $\beta$ -lactamase producing multidrug resistant *Klebsiella pneumoniae* isolates from pediatrics. *Pakistan J Zool* **2019**; 51(4): 1251–1257.

43. Tiwari B, Drogui P, Tyagi RD. Multidrug-resistant genes and pathogenic bacteria in hospital wastewater. In: Tyagi RD, Sellamuthu B, Pandey A, editors. *Current developments in biotechnology and bioengineering: environmental and health impact of hospital wastewater*; 2020. pp. 177–202.
44. Isturiz R. Global resistance trends and the potential impact on empirical therapy. *Int J Antimicrob Agents* 2008; 32: S201–S206.
45. Ju F, Beck K, Yin X, Maccagnan A, McArdell CS, Singer HP, et al. Wastewater treatment plant resistomes are shaped by bacterial composition, genetic exchange, and upregulated expression in the effluent microbiomes. *ISME J* 2019; 13(2): 346–360.
46. Dayan N, Dabbah H, Weissman I, Aga I, Even L, Glikman D. Urinary tract infections caused by community-acquired extended-spectrum  $\beta$ -lactamase-producing and nonproducing bacteria: a comparative study. *J Pediatr* 2013; 163(5): 1417–1421.

**How to cite this article:** Iqbal Z, Mumtaz MZ, Malik A. Extensive drug-resistance in strains of *Escherichia coli* and *Klebsiella pneumoniae* isolated from paediatric urinary tract infections. *J Taibah Univ Med Sc* 2021;16(4):565–574.