

Analysis of soft gelatin capsule with real-time polymerase chain reaction for halal authentication

Nina Salamah^{1,2*}, Any Guntarti^{1,2}, Laela Hayu Nurani¹

¹Faculty of Pharmacy, Universitas Ahmad Dahlan,

Jl. Prof. Dr. Soepomo, Janturan, Warungboto, Umbulharjo, Yogyakarta, Indonesia

²Ahmad Dahlan Halal Center, Universitas Ahmad Dahlan,

Jl. Prof. Dr. Soepomo, Janturan, Warungboto, Umbulharjo, Yogyakarta, Indonesia

Submitted: 24-01-2023

Reviewed: 06-03-2023

Accepted: 24-03-2023

ABSTRACT

Halal medicine is an interesting topic to always discuss because it is a priority choice for Muslim consumers, one of which is halal capsules. Currently, molecular biology techniques such as real-time polymerase chain reactions are rapidly developing, including for the analysis of non-halal components based on DNA sequences. This study aimed to validate the quantitative PCR method for identifying DNA in gelatin-based products and to apply the confirmation method designed for capsule samples on the market circulating in Yogyakarta to prove the halalness of these samples. Validation of the porcine DNA detection analysis method on standard extraction of porcine gelatin using primer pairs obtained in previous studies. Validated methods are used for testing market capsule shells. The qPCR method using D-loop primers is specifically capable of amplifying porcine gelatin DNA up to a concentration of 0.5 pg/ μ L, with a CV value in the amplification response of porcine gelatin DNA isolates (1000 pg/ μ L) of 0.85% which meets the test criteria using the PCR. Three samples of commercial soft capsules tested gave a positive amplification response, meaning that the samples tested contained porcine DNA, and one negative sample, which probably had non-porcine gelatin. The application of this method is also very useful for ensuring the authenticity of the capsule shell, especially from cross-contamination and counterfeiting.

Keywords: DNA, halal, porcine gelatin, real-time PCR, soft capsule

*Corresponding author:

Nina Salamah

Faculty of Pharmacy, Universitas Ahmad Dahlan

Jl. Prof. Dr. Soepomo, Warungboto, Umbulharjo, Yogyakarta, Indonesia

Email: nina.salamah@pharm.uad.ac.id



INTRODUCTION

The government creates a halal certificate instrument to guarantee the halalness of a product to protect consumers from the circulation of non-halal products (Hasan, 2014). This is due to the fact that not all goods available on the market are assured to be halal. As a result, business players are required by Article 4 of Law No. 33 of 2014 Concerning Halal Product Guarantee to possess a halal certificate and a halal logo (Anonymous, 2014). Products not including the halal logo and whose halal status is doubtful need to undergo a halal authentication process before being declared to contain non-halal ingredients. Therefore, product composition analysis regarding the presence or absence of non-halal components is necessary. The prospect of halal products, especially food, is developing rapidly. Indonesians consume halal products, reaching 1.8 billion people, and the current halal food market is predicted to be more than US 661 billion (Ali et al., 2017). Halal food is thought to be more widely known, and halal products are anticipated to grow rapidly (Rahman et al., 2014).

Approximately 10% of pharmaceutical preparations circulating in the community are capsules with gelatin's the main ingredient. Capsules are used for medicinal and preventive purposes, so most people must consume capsules. Capsule dosage forms provide various conveniences compared to other solid dosage forms of pharmaceutical products. Hence, manufacturers prefer to produce pork, beef, fish skin, and bone capsules to meet potential market needs. Therefore, a sensitive and specific method for confirming the origin of gelatin is needed.

Gelatin, a protein that can be obtained from the skin, connective tissue, and bones of agricultural animals like fish, poultry, cattle, and porcine, is created through the partial hydrolysis of collagen tissue. Gelatin from porcine is the most widely used in the medicine and food industry, considering that more gelatin from porcine is produced internationally; this is because it is easier for raw materials from porcine slaughter to be very large in quantity and ready to be used to create porcine gelatin compared to beef. The use of gelatin is increasingly diverse, but this needs to be matched by sufficient growth in domestic gelatin production so that until now, domestic needs have been met using imports. Based on data from the Central Statistics Agency (BPS) for 2020, the amount of gelatin imported into Indonesia reached 4808 tons worth 355 billion rupiahs. This statistical report shows how large the use of gelatin is in Indonesia, so it needs supervision regarding the certainty of halal gelatin (BPS, 2020).

Various kinds of gelatin and food products, both from imports and domestic industries, require clarity on the raw materials used, so the certainty of methods for detecting the raw materials used is urgently needed. Because non-halal ingredients are typically added to or mixed with biochemically similar matrices, the analysis of non-halal ingredients like porcine gelatin and pork in food items can be complicated. For instance, meatballs with (non-halal) pork incorporated are marketed as beef meatballs (Rohman et al., 2017), and porcine gelatin may be used in the manufacture of capsule shells, which are usually prepared from beef gelatin.

This study aimed to validate the quantitative PCR method for identifying DNA in gelatin-based products and to apply the confirmation method designed for samples of gelatin-based products on the market circulating in Yogyakarta to prove the halalness of these samples. In this study, the challenge was extracting DNA from modified capsule products to produce better DNA quality. This study is urgent because it uses qPCR to analyze porcine DNA to identify it and determine its halal status in a variety of goods. The negative results from this study still need to be confirmed using specific primers for bovine DNA to prove that the absence of porcine DNA indicates that bovine DNA is present in the product. The application of this method is also very useful for ensuring the authenticity of the capsule shell, especially from cross-contamination and counterfeiting.

MATERIALS AND METHODS

Primer design

Primer design was carried out using the online software PrimerQuest Tool with access code AF034253 for DNA DLOOP. The primer design aimed to obtain a specific SS2 primer pair for the porcine DLOOP (*Sus scrofa*) sequence. The primer was chosen by considering several parameters of the primer design, namely having a length of 15-30 bp, 40-60% GC content, G and C nucleotides evenly

Analysis of halal ... (Salamah, et al.)

distributed along the primer and primer T_m ranging from 50-60°C (Borah, 2011). Based on the *in silico* design of DNA primers, a pair of porcine primers (SS2) were obtained which were used in this study, namely forward: 5'-ACTTCATGGAAGTTCATGATCCG-3', and reverse 5'-ATGTACGTTATGTCCCGTAACC-3'. This porcine primer has melting temperature characteristics of 58.20°C (F) and 57.95°C (R), GC content of 45.45% (F) and 40.00% (R) with an amplicon length of 103 base pairs (bp) (Salamah et al., 2021).

Preparation and isolation of DNA from gelatin and commercial capsule shell samples

The blended capsule shells were weighed approximately 250 mg, added with 800 µL of 70% ethanol, vortexed for 30 minutes, then incubated in a water bath at 65°C for 30 minutes with vortex every 15 minutes. Then each was added 800 µL lysis buffer and 20 µL proteinase K and homogenized, then incubated in a water bath at 65°C for 30 minutes with a vortex every 15 minutes. Add 10 µL of RNase and incubate in a water bath at 38°C. Then, phenol and 0.5x volume of MCHA each was added and shaken with a shaker for 40 minutes. The mixture was centrifuged for 30 minutes at 14500 rpm at room temperature.

The supernatant was also moved to a fresh Eppendorf, 1x chloroform was added, and the mixture was agitated for 15 minutes. After centrifuging the mixture for 10 minutes at 14500 rpm, the supernatant was moved to a new Eppendorf, and 0.1x volume of 3M Na-acetate pH 7.6 was then added. Absolute cold ethanol in a 2x volume was introduced and overnight incubated at -80°C. They were spun at 15000 rpm for 5 minutes at 4 °C. For additional analysis, the leftover ethanol was air-dried in LAF for approximately 10 minutes, to which 40 L of TE buffer was then added (Sambrook et al., 1989).

Instrumental analysis using qPCR

qPCR analysis was carried out using a 20 µL reaction mixture, as written in Table 1, and the qPCR conditions followed the data in Table 2.

Table 1. qPCR reaction mixture using *Evagreen* with a total volume of 20 µL

Formula	Volume (µL)
Evagreen	10
Primer Reverse SSB (5µM)	1.0
Primer Forward SSB (5µM)	1.0
DNA (50 ng)	2.0
NFW	Variety
Total Volume	20

Table 2. Program the temperature of the qPCR reaction in one amplification cycle

Phase	Condition
Initial Denaturation	95°C, 30 seconds
Denaturation	95°C, 5 seconds
Annealing	52°C, 30 seconds
Extension	72°C, 10 seconds
Number of cycles	40 cycles

Primary specificity test

A primer pair specificity test was carried out using qPCR at the optimum attachment temperature to amplify 50 ng of bovine and porcine DNA. This test was also performed on DNA isolated from pure bovine and porcine gelatin (Bio-Rad, 2006; Ponchel, 2006).

Sensitivity test of the qPCR method on gelatin and capsule shells as a comparison

This was done by making eight DNA dilutions (1000; 200; 100; 10; 5; 1; 0.5; 0.01 pg/µL) from pure bovine gelatin or a comparison capsule shell with 100% bovine gelatin. Sensitivity tests were also

performed on all comparison capsule shell formulas at the optimum sticking temperature (Soares et al., 2013).

Repeatability test

The repeatability test took one point on the standard linear curve from the 100% bovine gelatin DNA dilution series. The repeatability test was then taken from the comparison capsule DNA (pork-cow gelatin mixture), the concentration of which was the same in the four replications.

Analysis of commercial capsule shell samples

The samples tested came from 4 different soft capsule manufacturers, namely Market samples 1, 2, 3, and 4. The method used is the same for testing comparator capsule shells, starting from specificity, sensitivity, and repeatability tests.

RESULT AND DISCUSSION

Gelatin is a polypeptide derived from collagen found in the skin, connective tissue, and bones of various species, including cattle, pigs, fish, and even insects (Abdelfadeel, 2012). The most dominant amino acids in the formation of the primary structure of gelatin are glycine, proline, and hydroxyproline, which are arranged repeatedly. In the process of gel formation, the structure of the gelatin undergoes a conformational change from the primary structure (random coil) to the secondary structure (α -helix) to return to the initial structure of collagen in the form of a triple- α -helix through the formation of intra- and inter-peptide hydrogen bonds, thus forming a woven/crosslink. This change in gelatin structure can complicate the DNA isolation step. Therefore in the gelatin DNA isolation technique, a preparation step is required by dissolving the gelatin in 1 mL of PBS-ethanol solution (1:1).

Phosphate Buffered Saline (PBS) used during the DNA extraction process aims to prevent changes in the random coil gelatin structure into a triple- α -helix because the salts contained in PBS cause ionic bonds to form. As a result, the possibility of forming hydrogen bonds is slight, so crosslinks do not occur. Gelatin is a lyophilic colloid that interacts strongly with the dispersion medium. It is very soluble in the dispersion medium, and gelatin is a colloid that dissolves easily in water to expand easily. This situation will complicate the isolation process. Therefore absolute ethanol is added to prevent the gelation of gelatin in water through a dehydration mechanism. Ethanol is a solvent with a lower polarity than water, resulting in competition in forming bonds between gelatin-water-ethanol. This situation causes the stability of the colloidal gelatin to be damaged so that the gelatin does not expand and precipitate. After the sample was dissolved entirely in the preparation solution, added 1 mL of absolute ethanol was centrifuged for 3 minutes at 13000 rpm to precipitate the gelatin, then the supernatant was discarded. The resulting gelatin precipitate is treated like a meat sample to isolate DNA, starting from the cell lysis stage to dissolving the DNA isolate in TE buffer (Erwanto et al., 2014).

The quantitative analysis of DNA isolates using a spectrophotometer (nanodrop) revealed that protein contamination was suggested by a ratio of A260/A280 that was less than 1.8. A ratio of greater than 1.8, however, denoted the existence of RNA contamination. The outcomes of the quantitative analysis demonstrated that the separation process involved insufficient washing, resulting in a less-than-pure DNA sample. According to the [Codex Alimentarius Commission \(2010\)](#), a PCR method's validity is impacted by high DNA purity because it permits optimum amplification.

The results of the initial stage were carried out by testing the specificity of the primer at the optimum annealing temperature obtained in the previous test, with a running time of 30 cycles (Ponchel, 2006). [Figure 1](#) shows that the specific primer where only the target DNA, porcine gelatin, gave a positive signal, while bovine gelatin was not amplified.

Cytochrome B primer pairs specific for porcine DNA at an attachment temperature of 52.0°C were then tested for specificity against DNA isolates from pure gelatin, both bovine and porcine gelatin. In this test, 50 ng of bovine DNA was also used as a positive control. The amplification response of DNA isolates is shown in [Figure 1](#). The resulting amplification only came from porcine gelatin DNA isolates (RFU = 50, Ct = 26.85, and Tm = 80.50), and bovine gelatin DNA isolates (positive control) did not

experience amplification. This shows that the primer D-LOOP is specific only for porcine gelatin DNA with an annealing temperature of 52.0°C.




Figure 1. Amplification curve (A) and melt curve analysis (B) of CYTBWB2 primer on porcine gelatin DNA and bovine gelatin comparator DNA

After obtaining a specific primer and optimum annealing temperature, it is followed by method validation consisting of a sensitivity, linearity, and repeatability test. The type of validation used is partial validation for quantitative methods. Method validation used several porcine gelatin DNA samples from the porcine gelatin DNA dilution series. DNA extracted from porcine gelatin with several types of composition, as well as DNA isolated from commercial soft capsules.




Figure 2. Amplification curve (A) and melt curve analysis (B) in the sensitivity test for porcine gelatin DNA using primer

Based on the results of the sensitivity test in Figure 2, the Limit of Detection (LOD) value of porcine gelatin DNA in this method is at a concentration of 0.5 pg/μL. The next stage is the linearity test, as shown in Figure 3 and Table 3. The linearity test was carried out by constructing a standard curve based on the amplification results of 7 porcine gelatin DNA dilution series. The standard curve describes the linear relationship between the \log^{10} concentration series of porcine gelatin DNA and C_q (cycle quantification) values. The standard curve of the serial amplification results of porcine gelatin DNA dilution using 7 points at 5000, 1000, 500, 100, 10, and 0.5 pg/μL. The LOD obtained in this validation was very good because it had a value of 0.5 pg/μL. Meanwhile, in the primer developed by Arini et al. (2018), the LOD obtained was still quite high, namely 48 pg/μL. Making this standard curve also simultaneously calculates the efficiency value of the PCR amplification process. The purity of the reagents used, inconsistencies in the small volume pipetting process, which leads to low accuracy, the

presence of inhibitors in the PCR master mix, and the ability of the test, which depends on the specificity of the primer and the length of the target sequence of Template DNA, can all have an impact on the efficiency value (Muhammed et al., 2015; Svec et al., 2015). An excessively high efficiency value suggests that the presence of an inhibitor during the reaction is interfering with the target DNA's ability to multiply each cycle. At the same time, an efficiency value that is too small indicates incomplete primer attachment, which results in exponentially less template amplification in each cycle.

Table 3. Linearity test results of porcine gelatin DNA

Concentration (pg/ μ L)	Cq
0,5	25,51
10	25,69
100	26,14
500	26,38
1000	26,43
5000	27,08

The linear regression equation was also obtained from the standard amplification reaction curve for porcine gelatin DNA isolates (Table 3 and Figure 3), with an R² value of 0.962, a slope of 0.475, and a y-intercept of 26.168, so that the curve meets the criteria of fairly good linearity. The resulting efficiency value (E) (99.2%), so that the efficiency value obtained meets the acceptance requirements in the range of 90-110% (Bio-Rad, 2006). Linearity test results on porcine gelatin DNA with various concentrations also gave good results. The results found are consistent with the linearity curve's requirements using the values in (Widyasari et al., 2015).




Figure 3. Standard curve of porcine gelatin DNA amplification results with various concentrations using primers

By calculating the average standard deviation (SD) and coefficient of variation (CV) values, the repeatability test—the next validation method—is designed to assess the repeatability of the amplification findings from bovine gelatin DNA at a concentration of 1000 pg/L. Repeatability expresses the degree to which findings from several measurements taken from homogeneous samples in a short period of time under predetermined circumstances are comparable. The amplification data obtained can be seen in Figure 4. The amplification response repeatability test on porcine gelatin DNA isolates at a concentration of 1000 pg/ μ L resulted in a coefficient of variation (CV) of 0.85%. The CV value meets the recommended criteria for using the PCR method, namely $CV \leq 25\%$ (CAC, 2010). The obtained CV value was 0.85%, lower than the highest CV value permitted for q-PCR quantitative assays in accordance with Codex Alimentarius Commission. The resulting Ct (cycle threshold) data, average Ct, standard deviation, and coefficient of variation are shown in Table 4 (CAC, 2010).




Figure 4. DNA repeatability test results isolated from porcine gelatin at a concentration of 1000 pg/μL; amplification curve (A) and melt peak curve (B)

Table 4. DNA test results for porcine gelatin

Sample	Cq	Average Cq	SD	CV (%)
Porcine gelatin DNA concentration 1000 pg/μL	27.19 26.87 27.16 26.54 26.92 27.11	26.97	0.23	0.85




Figure 5. The results of DNA isolates from soft capsule market samples, Amplification (A), and Melting curve analysis (B)

The studies obtained D-loop primers (forward: 5'-ACTTCATGGAACTCATGATCCG-3' and reverse 5'-ATGTACGTTATGTCCCGTAACC-3') that are specific for porcine gelatin DNA, with an attachment temperature of 52.0°C. The qPCR method can still provide an amplification response at a concentration of 0.5 pg/μL bovine DNA using D-loop primers, so this value is used as the LOD (Limit of Detection) value. Analysis of the repeatability of the amplification response from porcine gelatin DNA isolates gave a sequential CV value of 0.85%, which met the recommended CV value for the PCR method, namely $CV \leq 25\%$.

CONCLUSION

The qPCR method using D-loop primers is specifically capable of amplifying porcine gelatin DNA up to a concentration of 0.5 pg/ μ L, with a CV value in the amplification response of porcine gelatin DNA isolates (1000 pg/ μ L) of 0.85% which meets the test criteria using the PCR. Three samples of commercial soft capsules tested gave a positive amplification response, meaning that the samples tested contained porcine DNA, and one negative sample, which probably had non-porcine gelatin.

ACKNOWLEDGEMENT

The study was supported by regular research funds for 2022–2023, and the authors thank LPPM UAD for their support.

REFERENCES

- Abdelfadeel, H. F. A. (2012). *Extraction and characterization of gelatin from melon bug (Aspongubus viduatus) and sorghum bug (Agonoscelis pubescens) for application into ice-cream making*. Sudan University for Science & Technology.
- Ali, E., Sultana, S., Hamid, S. B. A., Hossain, M., Yehya, W. A., Kader, A., & Bhargava, S. K. (2017). Gelatin controversies in food, pharmaceuticals, and personal care products: Authentication methods, current status, and future challenges. *Critical Reviews in Food Science and Nutrition*, 58(9), 1495–1511. <https://doi.org/10.1080/10408398.2016.1264361>
- Anonymous. (2014). *Undang-Undang Nomor 33 Tahun 2014 tentang Jaminan Produk Halal*. Presiden Republik Indonesia. <https://peraturan.bpk.go.id/Home/Download/28038/UU%20Nomor%2033%20Tahun%202014.pdf>
- Arini, R. L., Ramadhani, D., Pebriyanti, N. W., Sismindari, S., & Rohman, A. (2018). The use of species-specific primer targeting on D-loop mitochondrial for identification of wild boar meat in meatball formulation. *Journal of Advanced Veterinary and Animal Research*, 5(3), 361. <https://doi.org/10.5455/javar.2018.e275>
- Badan Pusat Statistik (BPS). (2020). *Data Impor Produk Tertentu Tahun 2014-2019*. Badan Pusat Statistik.
- Bio-Rad. (2006). *Real Time PCR Application Guide*. Bio-Rad Laboratories, Inc.
- Borah, P. (2011). Primer designing for PCR. *Science Vision*, 11(3), 134–136.
- Codex Alimentarius Commission (CAC). (2010). *Guidelines on performance criteria and validation of methods for detection, identification, and quantification of specific DNA sequences and specific proteins in foods. CAC/GL 74-2010*. Codex Alimentarius Commission International Food Standards.
- Erwanto, Y., Abidin, M. Z., Muslim, E. Y. P., Sugiyono, S., & Rohman, A. (2014). Identification of pork contamination in meatballs of Indonesia local market using polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) analysis. *The Asian-Australian Journal Animal Sciences*, 27(10), 1487–1492. <https://doi.org/10.5713/ajas.2014.14014>
- Gelatin Manufacturers Institute of America (GMIA). (2012). *Gelatin Handbook*. Gelatin Manufacturers Institute of America. http://www.gelatin-gmia.com/uploads/1/1/8/4/118450438/gmia_gelatin_manual_2019.pdf
- Hasan, KN. S. (2014). Kepastian hukum sertifikasi dan labelisasi halal produk pangan. *Jurnal Dinamika Hukum*, 14(2), 227–238. <https://doi.org/10.20884/1.jdh.2014.14.2.292>
- Muhammed, M. A., Bindu, B. S. C., Jini, R., Prashanth, K. V. H., & Bhaskar, N. (2015). Evaluation of different DNA extraction methods for the detection of adulteration in raw and processed meat through polymerase chain reaction—restriction fragment length polymorphism (PCR-RFLP). *Journal of Food Science and Technology*, 52(1), 514–520. <https://doi.org/10.1007/s13197-013-1024-9>
- Ponchel, F. (2006). Real-time PCR using SYBR® Green. In M. T. Dorak (Ed.), *Real-Time PCR* (1st ed., p. 1). Taylor & Francis.
- Rahman, Md. M., Ali, Md. E., Hamid, S. B. A., Mustafa, S., Hashim, U., & Hanapi, U. K. (2014). Polymerase chain reaction assay targeting cytochrome b gene for the detection of dog meat

- adulteration in meatball formulation. *Meat Science*, 97(4), 404–409. <https://doi.org/10.1016/j.meatsci.2014.03.011>
- Rohman, A., Himawati, A., Triyana, K., Sismindari, S., & Fatimah, S. (2017). Identification of pork in beef meatballs using Fourier transform infrared spectrophotometry and real-time polymerase chain reaction. *International Journal of Food Properties*, 20(3), 654–661. <https://doi.org/10.1080/10942912.2016.1174940>
- Salamah, N., Erwanto, Y., Martono, S., & Rohman, A. (2021). The employment of real-time polymerase chain reaction using species-specific primer targeting on D-loop mitochondria for identification of porcine gelatin in soft candy. *Indonesian Journal of Chemistry*, 21(4), 852. <https://doi.org/10.22146/ijc.60413>
- Sambrook, J., Fritsch, E. R., & Maniatis, T. (1989). *Molecular Cloning A Laboratory Manual* (2nd ed.). Cold Spring Harbor Laboratory Press.
- Soares, S., Amaral, J. S., Oliveira, M. B. P. P., & Mafra, I. (2013). A SYBR Green real-time PCR assay to detect and quantify pork meat in processed poultry meat products. *Meat Science*, 94(1), 115–120. <https://doi.org/10.1016/j.meatsci.2012.12.012>
- Svec, D., Tichopad, A., Novosadova, V., Pfaffl, M. W., & Kubista, M. (2015). How good is a PCR efficiency estimate: Recommendations for precise and robust qPCR efficiency assessments. *Biomolecular Detection and Quantification*, 3, 9–16. <https://doi.org/10.1016/j.bdq.2015.01.005>
- Widyasari, Y. I., Sudjadi, S., & Rohman, A. (2015). Detection of rat meat adulteration in meat ball formulations employing real time PCR. *Asian Journal of Animal Sciences*, 9(6), 460–465. <https://doi.org/10.3923/ajas.2015.460.465>