

Stone recurrence among Indonesian kidney stone formers: a comprehensive analysis of genetic polymorphism, demographic, and clinical factors

Widi Atmoko^{1,2}, Missy Savira², Rosalina Fajriani², Sevita Sathya Wistara², Asmarinah³, Alida Roswita Harahap⁴, Fiastuti Witjaksono⁵, Retno Asti Werdhani⁶, Ferry Safriadi⁷, Agus Rizal Ardy Hariandy Hamid², Ponco Birowo², Nur Rasyid², Akmal Taher²



pISSN: 0853-1773 • eISSN: 2252-8083
<https://doi.org/10.13181/mji.oa.247596>
Med J Indones. 2024;33:190–7

Received: May 21, 2024

Accepted: October 03, 2024

Published online: November 08, 2024

Authors' affiliations:

¹Doctoral Program in Medical Sciences, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia, ²Department of Urology, Faculty of Medicine, Universitas Indonesia, Cipto Mangunkusumo Hospital, Jakarta, Indonesia,

³Department of Medical Biology, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia, ⁴Department of Clinical Pathology, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia,

⁵Department of Nutrition, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia, ⁶Department of Community Medicine, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia,

⁷Department of Urology, Faculty of Medicine, Universitas Padjadjaran, Hasan Sadikin Hospital, Bandung, Indonesia

Corresponding author:

Widi Atmoko

Department of Urology, Faculty of Medicine, Universitas Indonesia, Cipto Mangunkusumo Hospital, Jalan Salemba Raya No. 6, Central Jakarta 10430, DKI Jakarta, Indonesia

Tel/Fax: +62-21-3912477

E-mail: dr.widiatmoko@yahoo.com/
widi.atmoko01@ui.ac.id

ABSTRACT

BACKGROUND The recurrence rate of kidney stone disease (KSD) can exceed 50% within 5 years. This study aimed to investigate factors associated with stone recurrence based on single nucleotide polymorphisms (SNPs) of the calcium-sensing receptor (CaSR) rs1801725 and rs1042636, demographic, and clinical profiles.

METHODS We collected data on the demographic and clinical characteristics of 80 adult kidney stone formers from April 2021 to September 2022, including peripheral blood samples, morning and 24-hour urine, and 7-day water intake records. Real-time polymerase chain reaction was used to evaluate SNP. Comparisons were made between patients with recurrent and first-time stones.

RESULTS 41% of 80 patients have experienced stone recurrence. Recurrent stone formers exhibited significantly higher 24-hour urine calcium excretion ($p = 0.03$) and lower serum calcium levels ($p = 0.019$) than first-time stone formers. Hypocitraturia (100%), low urine volume (78%), and hyperoxaluria (55%) were the main abnormalities of all patients. No significant differences were found in CaSR gene polymorphisms and other demographic, biochemical, or clinical parameters.

CONCLUSIONS Recurrent stone formers had higher 24-hour urine calcium excretion and lower serum calcium levels. Other risk factors and CaSR polymorphisms may insignificantly affect KSD recurrence.

KEYWORDS genetic polymorphism, Indonesia, kidney stone, metabolic, recurrence, risk factors

Over the years, the incidence and burden of kidney stone disease (KSD) have increased globally.¹ The prevalence of KSD in Asia is approximately 1–19.1% and seems to be greater in Stone Belt regions, such as Southeast America, North Africa, Middle East and Southeast Asia, and the Northeast region of Australia, due to the impact of the hot climate on renal stone formation.² Moreover, the recurrence rate of KSD is

high, ranging from 0.9–94% within 5 years period.^{3,4} The pathophysiology of KSD involves a complex physiochemical process within the kidney.⁵ This process results from multiple causes, including genetic, biochemical, medical, socioeconomic, dietary habits or nutritional, and other environmental risk factors.⁶ A recently published meta-analysis of 53 articles on risk factors for KSD recurrence revealed that 12

risk factors were associated with stone recurrence. Nevertheless, the authors found that the follow-up period was heterogeneous, and publication bias existed for some risk factors, which may have affected the interpretability of the data.⁷

A twin study of genetic involvement in KSD found that the heritability of kidney stones is approximately 46–57%.⁸ Several studies have also found a correlation between single nucleotide polymorphisms (SNPs) and KSD,⁹ although the results in the literature have been inconsistent. Genes involved in calcium metabolism, such as the calcium-sensing receptor (CaSR), are among the most studied gene polymorphisms associated with KSD and are frequently observed in Asian populations.¹⁰ However, genetic polymorphisms reported to increase susceptibility or recurrence vary among different regions and ethnicities.^{11,12} A meta-analysis by Atmoko et al¹³ found that different polymorphisms were associated with an increased risk of kidney stone recurrence in Asians than in Caucasians. This underlines the fact that ethnicity and the multifactorial nature of KSD may explain the differences in genetic and SNP variants.¹³

Despite Indonesia being one of the most populous countries in the Stone Belt area, no study has thoroughly assessed of KSD recurrence. Many previously published articles from Indonesia were focused more on the surgical management of kidney stones. Therefore, a significant knowledge gap exists in the basic and clinical analyses of kidney stones. Improved preventive intervention strategies for patients can be developed with the assistance of risk factors for KSD recurrence. This study aimed to investigate the differences in CaSR variant rs1801725 and rs1042636 gene polymorphisms, demographic, and clinical profiles between first-time and recurrent patients with kidney stones in the Indonesian population. The findings of this study fill the gap in the literature and serve as a foundation for preventive measures against KSD recurrence.

METHODS

Study population

The study was conducted at the Cipto Mangunkusumo Hospital, Jakarta, Indonesia, between April 2021 and September 2022. The study protocol conformed to the Declaration of Helsinki and was approved by the Ethics Committee of the Faculty of

Medicine, Universitas Indonesia (No: KET-544/UN2.F1/ETIK/PPM.00.02/2020). The inclusion criteria included patients aged ≥ 18 years and meeting the diagnostic criteria for KSD based on ultrasonography, computed tomography (CT) scan, kidney, ureter, and bladder X-ray imaging. Those who met the inclusion criteria were consecutively recruited after providing informed consent. Patients were excluded if they refused to provide informed consent or had a horseshoe kidney, solitary kidney, medullary sponge kidney, polycystic kidney, congenital vesicoureteral reflux, neurogenic bladder, kidney cancer, pyonephrosis, a history of struvite stones, used urinary catheter or nephrostomy. This study included 80 patients who fulfilled the inclusion criteria and provided informed consent.

Baseline data, including body mass index (BMI), waist circumference (WC), and blood pressure, were retrieved during the screening process. All participants were Indonesian residents, and BMI was classified based on the Asia-Pacific cut-off criteria.¹⁴ High WC was defined as ≥ 80 cm for females and ≥ 90 cm for males.¹⁵ Patients with at least two symptomatic episodes or new stones after treatment were classified in the recurrence group, while the rest were classified into the first-time stone group. Education levels were classified as follows: high education level for those who completed senior high school or higher, intermediate level for those who completed junior high school, and low education level for those who completed only elementary school or had no formal education.

From each participant, an 8 ml of venous blood sample was collected for biochemical and genotyping analyses. Hypercalcemia, high serum creatinine levels, and hyperglycemia were defined as >10.2 mg/dl, >1.3 mg/dl, and >200 mg/dl, respectively. A cut-off of >6 mg/dl or >7.2 mg/dl was used for hyperuricemia in women and men, respectively. First-voided morning urine samples were collected to measure inflammatory cytokine gene expression. They were also required to collect their 24-hour urine samples using the S-Y Easy Fold 24hr Urine Collector (Shih-Yung Medical Instrument Co., Taiwan) without additional preservatives and store them in the refrigerator. The cut-off values for interpretation were $<2,000$ ml/day for low urine volume, hypercalciuria >250 mg/day for females and >300 mg/day for males, hyperoxaluria >40 mg/day, hyperuricosuria >750 mg/day, hypomagnesuria <60 mg/day, and hypocitraturia <300 mg/day.^{16,17}

Finally, all participants were asked to complete a validated self-administered 7-day fluid record (Liq.In7) to evaluate total daily water and beverage intake.¹⁸ All data were collected and managed using the Research Electronic Data Capture (REDCap®, USA) database hosted at Universitas Indonesia.¹⁹

Biochemical analyses

Blood samples were analyzed to determine random blood glucose (RBG), serum uric acid, and serum calcium levels using an Abbott Architect Chemistry Analyzer (Abbott Diagnostics, USA). Creatinine levels were assessed using isotope dilution gas chromatography-mass spectrometry. Urine pH was determined using a pH paper strip, with acidic urine defined as a pH <5.5. The 24-hour urine excretion of calcium, oxalate, uric acid, magnesium, and citrate was estimated using a colorimetric assay kit (BioVision, USA).

Genotyping

Genomic DNA was extracted from whole blood samples using the QIAamp DNA Blood Mini Kit (QIAGEN, Netherlands). The quantity and quality of the extracted DNA were measured at 260 and 280 nm using a NanoDrop spectrophotometer (Thermo Fisher Scientific, USA). Genotyping of SNPs was performed using the 7500 Fast and 7500 Real-Time PCR Systems (Thermo Fisher Scientific). TaqMan GTXpress Master Mix and SNP genotyping assays (Thermo Fisher Scientific) were used to detect alleles present in the samples.

Statistical analysis

Data were presented as means and standard deviations, or medians and interquartile ranges for continuous variables, whereas categorical variables were presented as proportions. Statistical comparisons were performed using an independent t-test or the Mann-Whitney U test for continuous variables. All data analyses were completed using SPSS software version 25.0 (IBM Corp., USA), with $p < 0.05$ indicating statistical significance. Genotypic, dominant, and recessive genotyping models were used to analyze the association of genes with KSD recurrence.

RESULTS

Demographic and clinical characteristics

This study examined 80 patients, including 33 (41%) with recurrent stones and 47 (59%) with first-

time stones. The patients' ages ranged from 46 to 63 years, with a median age of 55 years. The proportions of office employees, shift workers, individuals with multiple stones, individuals with a family history of KSD, individuals with overweight obesity, and individuals with a large WC were higher among recurrent stone formers than among first-time stone formers. However, none of the differences in these proportions were statistically significant (Table 1). Average water consumption was low, with a mean of approximately 1,837 (539) ml for all participants.

Biochemical characteristics

Recurrent stone formers exhibited significantly elevated 24-hour urine calcium excretion ($p = 0.03$), although their serum calcium levels were low ($p = 0.019$). They also presented higher RBG levels, 24-hour urine oxalate, uric acid, citrate excretion, and magnesium excretion. The main abnormalities found in all participants were hypocitraturia (100%), low urine volume (78%), and hyperoxaluria (55%), with comparable proportions of single-type abnormalities between the groups. The results of the biochemical analysis are shown in Table 1.

Gene polymorphism and kidney stone recurrence

We investigated SNPs in the CaSR variants rs1801725 and rs1042636 genes, and their distributions were comparable between recurrent and first-time stone formers based on genotyping, dominant, and recessive models (Table 2).

DISCUSSION

Without preventive measures, the relapse rate of KSD is estimated to be 10–23% per year, 50% within 5–10 years, and 75% within 20 years after the first occurrence.¹⁵ In this study, 41% of the participants had recurring stones, which is consistent with the rate reported in the literature.⁷ A good proportion of patients with KSD recurrence in this study can be rationalized, as this study was mostly conducted on patients with complex cases in a national referral hospital.

This study aligns with a previous study that failed to find any association between sex and KSD recurrence,²⁰ despite the finding that patients with recurring and first-time stones were predominantly male. In contrast, Daudon et al²¹ found that KSD recurrence was more common in men, based on an analysis of almost

Table 1. Demographic, clinical, and biochemical features of the subjects

Characteristics	Recurrent stone former (N = 33)	First-time stone former (N = 47)	p
Demographic and clinical features			
Sex, n (%)			0.272
Male	17 (52)	30 (64)	
Female	16 (48)	17 (36)	
Age (years), median (IQR)	55.0 (46.5–63.0)	55.0 (46.0–60.0)	0.604*
Degree of education, n (%)			
High	18 (55)	20 (43)	0.578
Intermediate	10 (30)	19 (40)	0.804
Low	5 (15)	8 (17)	0.557
Occupation, n (%)			0.883
Office employees	16 (48)	22 (47)	
Non-office employee	17 (52)	25 (53)	
Shift worker, n (%)			1.000
Yes	1 (3)	0 (0)	
No	32 (97)	47 (100)	
Multiple stones, n (%)			0.141
Yes	23 (70)	25 (53)	
No	10 (30)	22 (47)	
Family history of KSD, n (%)			0.089
Yes	12 (36)	9 (19)	
No	21 (64)	38 (81)	
BMI (kg/m ²), median (IQR)	26.3 (23.15–29.5)	26.7 (22.3–28.9)	0.856*
BMI classification, n (%)			0.513
Overweight–obese	26 (79)	34 (72)	
Underweight–normal	7 (21)	13 (28)	
WC (cm), mean (SD)	96.91 (13.3)	95.17 (12.39)	0.681
WC classification, n (%)			0.931
High	25 (76)	36 (77)	
Normal	8 (24)	11 (23)	
Daily water intake (ml), mean (SD)	1,866 (541)	1,817 (543)	0.694
Biochemical features			
Serum calcium (mg/dl), mean (SD)	8.77 (0.55)	9.14 (0.77)	0.019
High, n (%)	0 (0)	2 (4)	0.999
Serum uric acid (mg/dl), median (IQR)	6.4 (4.9–7.95)	6.7 (6.0–8.6)	0.122*
High, n (%)	13 (39)	25 (53)	0.225
Serum creatinine (mg/dl), median (IQR)	1.0 (0.9–1.4)	1.1 (0.75–1.55)	0.453*
High, n (%)	10 (30)	14 (30)	0.960
RBG (mg/dl), median (IQR)	102 (91.5–149.5)	99 (81–134.0)	0.214*
High, n (%)	5 (15)	2 (4)	0.110
24-hour urine volume (ml), median (IQR)	1,450 (975–2,025)	1,300 (950–1,800)	0.577*
High, n (%)	24 (73)	38 (81)	0.394
Urine calcium (mg/24 hour), median (IQR)	101.2 (52.96–168.88)	62.09 (41.85–107.39)	0.03*
High, n (%)	2 (6)	0 (0)	0.999

Continued on next page

Table 1. (Continued)

Characteristics	Recurrent stone former (N = 33)	First-time stone former (N = 47)	<i>p</i>
Urine oxalate (mg/24 hour), median (IQR)	65.53 (33.45–165.7)	48.63 (25.5–105.71)	0.276*
High, n (%)	18 (55)	26 (55)	0.945
Urine uric acid (mg/24 hour), median (IQR)	49.27 (26.12–75.83)	37.31 (24.27–72.67)	0.639*
High, n (%)	0 (0)	0 (0)	NA
Urine magnesium (mg/24 hour), median (IQR)	65.47 (43.61–92.58)	79.7 (45.48–109.35)	0.195*
High, n (%)	15 (45)	14 (30)	0.154
Urine citrate (mg/24 hour), median (IQR)	11.59 (7.49–19.73)	8.15 (2.41–18.21)	0.121*
High, n (%)	33 (100)	47 (100)	NA
Urine pH, median (IQR)	6.0 (5.25–6.5)	6.0 (6.0–6.0)	0.917*
High, n (%)	8 (24)	9 (19)	0.584
Abnormal biochemical indices			
UTI, n (%)			0.536
High	9 (27)	10 (21)	

BMI=body mass index; IQR=interquartile range; KSD=kidney stone disease; NA=not available; RBC=random blood glucose; SD=standard deviation; UTI=urinary tract infection; WC=waist circumference

*Mann-Whitney test

The cut-off values for interpretation were >10.2 mg/dl for hypercalcemia; >750 mg/day for hyperuricosuria; >1.3 mg/dl for high serum creatinine levels; >200 mg/dl for hyperglycemia; <2,000 ml/day for low urine volume; >250 mg/day or >300 mg/day for hypercalciuria in women and men, respectively; >40 mg/day for hyperoxaluria; >6 mg/dl or >7.2 mg/dl for hyperuricemia in women and men, respectively; <60 mg/day for hypomagnesuria; <300 mg/day for hypocitraturia; and pH <5.5 for acidic urine

17,000 participants. It was previously hypothesized that estrogen in women might reduce calcium receptor expression, crystal binding capacity, and adenosine triphosphate in renal tubular cells, creating a protective environment against KSD.²² Nevertheless, temporal data have shown increased KSD incidence among females in recent years, particularly among adolescents.²³

This study had a median age of 55 years, supporting the finding of Widyasmara et al²⁴ that the peak incidence of kidney stones in Indonesia occurs between 51 and 60 years. Moreover, Trinchieri et al²⁰ reported no age differences between patients with one stone recurrence and those with first-time stones, which contrasts with a study by Li et al,²⁵ who reported that patients aged <20 years had higher rates of multiple stone recurrences. This study did not include patients aged <18 years, which may explain why the ages of the patients in the two groups were comparable. Moreover, no age data on the beginning of kidney stone formation was available; therefore, patients in the recurrent group may have had their first kidney stone occurring at a much younger age.

In this study, there were more participants with intermediate-to-higher education levels than those with low education levels in both groups. Additionally,

more than 40% of the participants were office workers. Sedentary occupations lead to a higher risk of kidney stones, particularly for those with low water intake habits due to limited access to water or sanitary toilets.²⁶ Nevertheless, based on our findings, neither educational status nor occupation correlated with KSD recurrence.

This study suggests that a family history of KSD and multiple renal stones were more common in the recurrence group, although the difference was not statistically significant. This finding aligns with some related research^{20,27} but contrasts with the meta-analysis findings by Wang et al⁷ in 2022, which identified family history as a significant recurrence risk. Recurrence intervals were notably shorter in patients with a family history.⁷ One study also showed that patients with multiple stones had a greater risk of recurrence than those with a single stone.²⁸ The difference in the results may stem from the smaller sample size compared to earlier studies.

Clinical factors, particularly hydration levels, can affect kidney stone recurrence. A previous study in Indonesia showed that most adults met their fluid intake recommendations, averaging over 2,000 ml/day.²⁹ However, the patients with kidney stones in our study only consumed approximately 1,837 (539) ml

Table 2. Association of CaSR SNP genotype to stone recurrence

SNP	Model	Genotype	Recurrent stone former, n (%) (N = 33)	First-time stone former, n (%) (N = 47)	p
CaSR rs1801725	Genotypic	TT	0 (0)	2 (4)	0.999
		GT	5 (15)	12 (26)	0.229
		GG	28 (85)	33 (70)	1.00
	Dominant	GT + TT	5 (15)	14 (29.8)	0.136
		GG	28 (85)	32 (70.2)	1.00
	Recessive	TT	0 (0)	2 (4)	0.999
GG + GT		33 (100)	45 (96)	1.00	
CaSR rs1042636	Genotypic	GG	10 (30)	14 (30)	0.355
		AG	18 (55)	20 (43)	0.169
		AA	5 (15)	13 (28)	1.00
	Dominant	AG + GG	28 (85)	34 (72)	0.269
		AA	5 (15)	13 (28)	1.00
	Recessive	GG	10 (30)	14 (30)	0.960
AA + AG		23 (70)	33 (70)	1.00	

CaSR=calcium-sensing receptor; SNP=single nucleotide polymorphism

daily, leading to a low urine volume. Compared with first-time stone formers, recurrent stone formers had significantly greater daily water intake and 24-hour urine volume, supporting a previous study that found lower urine volume was more common in first-time stone formers.³⁰ A daily urine output of 2.5 liters is recommended to reduce KSD risk.³¹

Metabolic syndrome is also linked to kidney stone recurrence risk, with factors such as low urine pH and high excretion of oxalate, uric acid, and phosphate often observed in individuals with high BMI.³² Obesity also correlates with high stone recurrence risk.³³ However, this study found no significant differences in BMI or WC between the two groups. A chronic hyperglycemic state is associated with KSD formation, and the severity of diabetes is a key predictor of recurrence among patients with KSD and type 2 diabetes mellitus (DM).^{34,35} Insulin resistance in DM leads to decreased renal ammonium production and decreased urinary pH, thus creating a favorable environment for uric acid stone formation.³⁶

This study revealed that hypocitraturia, low urine volume, and hyperoxaluria were the most common abnormalities. Notably, 4% of our patients had hypercalcemia, 6% had hypercalciuria, and none had hyperuricosuria. We agree with a recent systematic review of 28 articles that concluded hyperoxaluria and hypocitraturia are common in non-Western countries.³⁷

In contrast, a higher incidence of hypercalciuria is found in Western countries.³⁷ These differences in disease risk may be attributed to various dietary patterns and nutritional compositions, and ethnicity may also contribute to these differences in disease risk.³⁸

Despite a lower rate of hypercalciuria, we found that higher urine calcium excretion was significantly and independently associated with KSD recurrence, which is consistent with the findings of Hong et al.³⁹ According to the univariate analysis, patients in the recurrent stone group also had lower serum calcium levels, which could be associated with KSD if coupled with calcium loss in the kidney. Despite this discovery, the recurrent and first-time stone formation groups showed similar rates of hypercalciuria and hypercalcemia. Therefore, although urine calcium excretion levels did not fall in the hypercalciuria group, patients with high-normal levels should be cautioned about the possibility of stone recurrence. Urine supersaturation resulting from calcium excretion puts patients at a risk of stone formation. This is important because a previous study revealed that calcium oxalate is the most common stone component in the Indonesian population, followed by uric acid.²⁴

Hypomagnesuria is also significantly associated with kidney stone recurrence. Magnesium prevents KSD by inhibiting calcium oxalate crystallization, which

is enhanced when combined with citrate and acidic urine.⁴⁰ Increased dietary magnesium intake is also associated with a reduced risk of KSD.⁴¹ Future studies are needed to determine the role of magnesium supplementation in KSD recurrence.

Although 48% of our patients had hyperuricemia, none had hyperuricosuria. Similarly, several previous studies have shown that hyperuricemia is not associated with hyperuricosuria.^{42,43} Therefore, hyperuricemia should not be used as a predictive factor for hyperuricosuria, considering that patients with normal serum urate levels may be overlooked during treatment. However, the evaluation of serum urate concentration still needs to be considered in patients with a history of arthropathy or gout.

CaSR gene, located in chromosome 3q13.3–21, can express the tubular cells protein in thick ascending limbs to regulate calcium reabsorption by inhibiting paracellular uptake of calcium, resulting in hypercalciuria.⁴⁴ A previous study found that a variant of the minor T allele in CaSR rs1801725 and the minor G allele in rs1042636 significantly increased the risk of KSD in Eastern India and the Egyptian population.^{11,45} Furthermore, a combination of GT + TT genotypes in rs1801725 results in significantly higher serum calcium levels.¹¹ This result agrees with several other studies that found the association of a minor T allele in SNP rs1801725 to serum calcium level.⁴⁶ In contrast, rs1801725 was not correlated with an increased risk of KSD among the Chinese population.⁴⁷ This study has found no significant correlations between CaSR (rs1801725 and rs1042636) and kidney stone recurrence using any models. This study is the first to analyze CaSR SNPs associated with KSD recurrence in Indonesia. The small sample size, lack of healthy controls, and different genetic and phenotypic backgrounds may explain these findings. Therefore, it is premature to draw firm conclusions about the role of SNPs in KSD recurrence.

This study has some limitations. Due to a small patient cohort, ultrasound and kidney, ureter, and bladder X-ray, which are less sensitive than CT, were used. Other limitations include the lack of stone composition, comparative analysis with a control group, and failure to consider participant morbidities and medications. Future research should include case-control studies for a more comprehensive risk factor analysis to validate the current research findings.

Our study provides the groundwork for future studies on kidney stone pathophysiology, particularly

in the Indonesian context. This study has significant implications for many beneficiaries, including clinicians, patients, and policymakers. Clinicians and patients will benefit from better guidance on personalized and effective kidney stone prevention and treatment strategies. Policymakers can use these insights to develop public health initiatives to reduce kidney stone recurrence.

In conclusion, recurrent stone formers had higher 24-hour urine calcium excretion and lower serum calcium levels. An insignificantly higher proportion of risk factors, such as occupation type, family history of KSD, and obesity, was found; however, no difference was found in CaSR polymorphisms (rs1801725 and rs1042636) between the recurrent and first-time stone groups. Given the small sample size and the complexity of KSD recurrence, this finding cannot be generalized to larger or different populations. Further studies with larger sample sizes are needed.

Conflict of Interest

Agus Rizal Ardy Hariandy Hamid is the editor-in-chief of this journal but was not involved in the review or decision-making process of the article.

Acknowledgment

We thank all the subjects who participated in this study.

Funding Sources

None.

REFERENCES

1. Stamatelou K, Goldfarb DS. Epidemiology of kidney stones. *Healthcare (Basel)*. 2023;11(3):424.
2. Liu Y, Chen Y, Liao B, Luo D, Wang K, Li H, et al. Epidemiology of urolithiasis in Asia. *Asian J Urol*. 2018;5(4):205–14.
3. Baowaidan F, Zugail AS, Lyoubi Y, Culty T, Lebdai S, Brassart E, et al. Incidence and risk factors for urolithiasis recurrence after endourological management of kidney stones: a retrospective single-centre study. *Prog Urol*. 2022;32(8–9):601–7.
4. Vaughan LE, Enders FT, Lieske JC, Pais VM, Rivera ME, Mehta RA, et al. Predictors of symptomatic kidney stone recurrence after the first and subsequent episodes. *Mayo Clin Proc*. 2019;94(2):202–10.
5. Tamborino F, Cicchetti R, Mascitti M, Litterio G, Orsini A, Ferretti S, et al. Pathophysiology and main molecular mechanisms of urinary stone formation and recurrence. *Int J Mol Sci*. 2024;25(5):3075.
6. Prezioso D, Strazzullo P, Lotti T, Bianchi G, Borghi L, Caione P, et al. Dietary treatment of urinary risk factors for renal stone formation. A review of CLU Working Group. *Arch Ital Urol Androl*. 2015;87(2):105–20. Erratum in: *Arch Ital Urol Androl*. 2016;88(1):76.
7. Wang K, Ge J, Han W, Wang D, Zhao Y, Shen Y, et al. Risk factors for kidney stone disease recurrence: a comprehensive meta-analysis. *BMC Urol*. 2022;22(1):62.
8. Goldfarb DS, Avery AR, Beara-Lasic L, Duncan GE, Goldberg J. A twin study of genetic influences on nephrolithiasis in women and men. *Kidney Int Rep*. 2018;4(4):535–40.

9. Vladimirovna FT, Faridovich KK, Igorevich RV, Mikhailovich RL, Georgievich TD, Victorovich ED, et al. Genetic factors of polygenic urolithiasis. *Urologia*. 2020;87(2):57–64.
10. Mohammadi A, Shabestari AN, Baghdadabab LZ, Khatami F, Reis LO, Pishkuhi MA, et al. Genetic polymorphisms and kidney stones around the globe: a systematic review and meta-analysis. *Front Genet*. 2022;13:913908.
11. Guha M, Bankura B, Ghosh S, Pattanayak AK, Ghosh S, Pal DK, et al. Polymorphisms in CaSR and CLDN14 genes associated with increased risk of kidney stone disease in patients from the eastern part of India. *PLoS One*. 2015;10(6):e0130790.
12. Chen G, Hu C, Song Y, Xiu M, Liang W, Ou N, et al. Relationship between the Apal (rs7975232), Bsm1 (rs1544410), FokI (rs2228570), and TaqI (rs731236) variants in the vitamin D receptor gene and urolithiasis susceptibility: an updated meta-analysis and trial sequential analysis. *Front Genet*. 2020;11:234.
13. Atmoko W, Raharja PA, Birowo P, Hamid AR, Taher A, Rasyid N. Genetic polymorphisms as prognostic factors for recurrent kidney stones: a systematic review and meta-analysis. *PLoS One*. 2021;16(5):e0251235.
14. Pan WH, Yeh WT. How to define obesity? evidence-based multiple action points for public awareness, screening, and treatment: an extension of Asian-Pacific recommendations. *Asia Pac J Clin Nutr*. 2008;17(3):370–4.
15. World Health Organization. Regional Office for the Western Pacific. The Asia-Pacific perspective: redefining obesity and its treatment [Internet]. Sydney: Health Communications Australia; 2000. Available from: <https://iris.who.int/handle/10665/206936>.
16. Ennis JL, Asplin JR. The role of the 24-h urine collection in the management of nephrolithiasis. *Int J Surg*. 2016;36(Pt D):633–37.
17. Kumar R, Kapoor R, Mittal B, Kumar A, Mittal RD. Evaluation of urinary abnormalities in urolithiasis patients: a study from North India. *Indian J Clin Biochem*. 2003;18(2):209–15.
18. Johnson EC, Péronnet F, Jansen LT, Capitan-Jiménez C, Adams JD, Guelinckx I, et al. Validation testing demonstrates efficacy of a 7-day fluid record to estimate daily water intake in adult men and women when compared with total body water turnover measurement. *J Nutr*. 2017;147(10):2001–7.
19. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377–81.
20. Trinchieri A, Ostini F, Nespoli R, Rovera F, Montanari E, Zanetti G. A prospective study of recurrence rate and risk factors for recurrence after a first renal stone. *J Urol*. 1999;162(1):27–30.
21. Daudon M, Jungers P, Bazin D, Williams JC Jr. Recurrence rates of urinary calculi according to stone composition and morphology. *Urolithiasis*. 2018;46(5):459–70.
22. Peerapen P, Thongboonkerd V. Protective cellular mechanism of estrogen against kidney stone formation: a proteomics approach and functional validation. *Proteomics*. 2019;19(19):e1900095.
23. Gillams K, Juliebø-Jones P, Juliebø SØ, Somani BK. Gender differences in kidney stone disease (KSD): findings from a systematic review. *Curr Urol Rep*. 2021;22(10):50.
24. Widiasmara HB, Birowo P, Rasyid N. Urinary stone composition analysis in Indonesian population: a single major centre analysis. *Indo J Urol*. 2018;25(2):104–9.
25. Li Y, Bayne D, Wiener S, Ahn J, Stoller M, Chi T. Stone formation in patients less than 20 years of age is associated with higher rates of stone recurrence: Results from the Registry for Stones of the Kidney and Ureter (ReSKU). *J Pediatr Urol*. 2020;16(3):373.e1–6.
26. Malieckal DA, Goldfarb DS. Occupational kidney stones. *Curr Opin Nephrol Hypertens*. 2020;29(2):232–6.
27. Koyuncu HH, Yencilek F, Eryildirim B, Sarica K. Family history in stone disease: how important is it for the onset of the disease and the incidence of recurrence? *Urol Res*. 2010;38(2):105–9.
28. Kang HW, Seo SP, Kwon WA, Woo SH, Kim WT, Kim YJ, et al. Distinct metabolic characteristics and risk of stone recurrence in patients with multiple stones at the first-time presentation. *Urology*. 2014;84(2):274–8.
29. Laksmi PW, Morin C, Gandy J, Moreno LA, Kavouras SA, Martinez H, et al. Fluid intake of children, adolescents and adults in Indonesia: results of the 2016 Liq.In⁷ national cross-sectional survey. *Eur J Nutr*. 2018;57(Suppl 3):89–100.
30. Abu-Ghanem Y, Shvero A, Kleinmann N, Winkler HZ, Zilberman DE. 24-h urine metabolic profile: is it necessary in all kidney stone formers? *Int Urol Nephrol*. 2018;50(7):1243–7.
31. Skolarikos A, Jung H, Neisius A, Petfik A, Somani B, Taily T, et al. EAU guidelines on urolithiasis [Internet]. Arnheim: European Association of Urology; 2023. Available from: <https://uroweb.org/guidelines/urolithiasis>.
32. Taylor EN, Curhan GC. Body size and 24-hour urine composition. *Am J Kidney Dis*. 2006;48(6):905–15.
33. Zeng J, Wang S, Zhong L, Huang Z, Zeng Y, Zheng D, et al. A retrospective study of kidney stone recurrence in adults. *J Clin Med Res*. 2019;11(3):208–12.
34. Geraghty R, Abdi A, Somani B, Cook P, Roderick P. Does chronic hyperglycaemia increase the risk of kidney stone disease? results from a systematic review and meta-analysis. *BMJ Open*. 2020;10(1):e032094.
35. Torricelli FC, De S, Gebreselassie S, Li I, Sarkissian C, Monga M. Type-2 diabetes and kidney stones: impact of diabetes medications and glycemic control. *Urology*. 2014;84(3):544–8.
36. Nerli R, Jali M, Guntaka AK, Patne P, Patil S, Hiremath MB. Type 2 diabetes mellitus and renal stones. *Adv Biomed Res*. 2015;4:180.
37. Huynh LM, Dianatnejad S, Tofani S, Carrillo Ceja R, Liang K, Tapiero S, et al. Metabolic diagnoses of recurrent stone formers: temporal, geographic and gender differences. *Scand J Urol*. 2020;54(6):456–62.
38. Donin AS, Nightingale CM, Owen CG, Rudnicka AR, McNamara MC, Prynne CJ, et al. Nutritional composition of the diets of South Asian, black African-Caribbean and white European children in the United Kingdom: the child heart and health study in England (CHASE). *Br J Nutr*. 2010;104(2):276–85.
39. Hong YH, Dublin N, Razack AH, Mohd MA, Husain R. Urinary metabolic evaluation of stone formers—a Malaysian perspective. *Urology*. 2012;80(3):529–34.
40. Riley JM, Kim H, Averch TD, Kim HJ. Effect of magnesium on calcium and oxalate ion binding. *J Endourol*. 2013;27(12):1487–92.
41. Shringi S, Raker CA, Tang J. Dietary magnesium intake and kidney stone: the national health and nutrition examination survey 2011–2018. *R I Med J* (2013). 2023;106(11):20–5.
42. Eyre KS, Lewis F, Cui H, Grout E, Mihai R, Turney BW, et al. Utility of blood tests in screening for metabolic disorders in kidney stone disease. *BJU Int*. 2021;127(5):538–43.
43. Sila-On A, Pavaro U, Nuchpramool W. Serum and urinary uric acid levels in healthy subjects and in patients with urolithiasis. *J Med Assoc Thai*. 1991;74(8):352–7.
44. Hendy GN, Canaff L. Calcium-sensing receptor gene: regulation of expression. *Front Physiol*. 2016;7:394.
45. Ali FT, Abd El-Azeem EM, Hekal HF, El-Gizawy MM, Sayed MS, Mandoh AY, et al. Association of TRPV5, CASR, and CALCR genetic variants with kidney stone disease susceptibility in Egyptians through main effects and gene–gene interactions. *Urolithiasis*. 2022;50:701–10.
46. Chang X, Li J, Guo Y, Wei Z, Mentch FD, Hou C, et al. Genome-wide association study of serum minerals levels in children of different ethnic background. *PLoS One*. 2015;10(4):e0123499.
47. Ding Q, Fan B, Shi Y, Fan Z, Ding L, Li F, et al. Calcium-sensing receptor genetic polymorphisms and risk of developing nephrolithiasis in a Chinese population. *Urol Int*. 2017;99(3):331–7.